

For Research Use Only

CoraLite® Plus 647-conjugated TNF Alpha Monoclonal antibody



Catalog Number: CL647-69002

Basic Information

Catalog Number: CL647-69002	GenBank Accession Number: GeneID (NCBI): 7124	Purification Method: Protein G purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	Full Name: tumor necrosis factor (TNF superfamily, member 2)	CloneNo.: 11H8A5
Source: Mouse		Excitation/Emission maxima wavelengths: 650 nm / 665 nm
Isotype: IgG1		
Immunogen Catalog Number: HZ-1014		

Applications

Tested Applications:
FC (Intra)

Species Specificity:
Human

Background Information

TNF, as also known as TNF-alpha, or cachectin, is a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. It is expressed as a 26 kDa membrane bound protein and is then cleaved by TNF-alpha converting enzyme (TACE) to release the soluble 17 kDa monomer, which forms homotrimers in circulation. It is produced chiefly by activated macrophages, although it can be produced by many other cell types such as CD4+ lymphocytes, NK cells, neutrophils, mast cells, eosinophils, and neurons. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, INS resistance, and cancer.

Storage

Storage:
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.




Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ PMA+ionomycin-stimulated and Monensin Solution treated human PBMCs were intracellularly stained with 0.25 ug CoraLite® Plus 647 Anti-Human TNF Alpha (CL647-69002, Clone:11H8A5), and 0.25 ug CoraLite®488 Anti-Human CD3 (UCHT1) (CL488-65151, Clone: UCHT1). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

1X10⁶ human PBMCs were intracellularly stained with 0.25 ug CoraLite® Plus 647 Anti-Human TNF Alpha (CL647-69002, Clone:11H8A5), and 0.25 ug CoraLite®488 Anti-Human CD3 (UCHT1) (CL488-65151, Clone: UCHT1). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

1X10⁶ NK92 were intracellularly stained with 0.25 ug CoraLite® Plus 647 Anti-Human TNF Alpha (CL647-69002, Clone:11H8A5) (red), or 0.25 ug x. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).