

For Research Use Only

CoraLite® Plus 647-conjugated Noggin Monoclonal antibody



Catalog Number:CL647-69010

Basic Information

Catalog Number: CL647-69010	GenBank Accession Number: GeneID (NCBI): 9241	Purification Method: Protein G purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	Full Name: noggin	CloneNo.: 2A11C1
Source: Mouse		Excitation/Emission maxima wavelengths: 650 nm / 665 nm
Isotype: IgG1		
Immunogen Catalog Number: HZ-1118		

Applications

Tested Applications:
FC (Intra)

Species Specificity:
Human

Background Information

Noggin is an extracellular polypeptide acting as an antagonist of bone morphogenetic proteins (BMPs) regulating embryonal development. Noggin inhibits activity of BMP-2, -4, -7, -13, and -14. Noggin is present extracellularly in the matrix or retained at the cell surface via interaction with heparin sulfate proteoglycans. In early development stages, Noggin is produced by the Spemann organizer, allowing dorsal-ventral patterning of BMPs (PMID: 8752214). Subsequently, Noggin is expressed by the notochord regulating BMP-4 signaling in neurogenesis. Additionally, Noggin is present during development in the dermal papilla, connective tissue of the hair follicle, lens, retina, and pericocular mesenchyme, as well as in the mesoderm lineage regulating development of the bone, cartilage, and muscles.

This antibody is used to Block/Neutralize Noggin protein.

Storage

Storage:
Store at -20°C. Avoid exposure to light.

Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ PC-3 cells were intracellularly stained with 0.8 ug CoraLite® Plus 647 Anti-Human Noggin (CL647-69010, Clone:2A11C1) (red), or 0.8 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).