## For Research Use Only

## CoraLite® Plus 647-conjugated Phospho-JUN (Ser73) Recombinant antibody



Catalog Number: CL647-80086

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL647-80086 BC068522 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 3725

Nanodrop: **UNIPROT ID:** 

Source: P05412 Rabbit Full Name:

Isotype: jun oncogene IgG

Calculated MW: 331 aa, 36 kDa Observed MW:

42-45 kDa

**Purification Method:** 

Protein A purification

CloneNo.: 4A18

Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity:

human, mouse

**Background Information** 

JUN, the most extensively studied protein of the activator protein-1 (AP-1) complex, is involved in numerous cell activities, such as proliferation, apoptosis, survival, tumorigenesis and tissue morphogenesis (PMID: 22180088). JUN is a transcription factor that recognizes and binds to the enhancer heptamer motif 5'-TGA[CG]TCA-3'. It promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation. JUN is a basic leucine zipper (bZIP) transcription factor that acts as homo- or heterodimer, binding to DNA and regulating gene transcription (PMID: 9732876). In additon, extracellular signals can induce post-translational modifications of JUN, resulting in altered transcriptional activity and target gene expression (PMID:8464713). More over, it has uncovered multiple layers of a complex regulatory scheme in which JUN is able to crosstalk, amplify and integrate different signals for tissue development and disease. Jun is predominantly nuclear, ubiquitinated Jun colocalizes with lysosomal proteins (PMID: 15469925). This antibody is raised against synthetic phosphopeptide corresponding to residues surrounding Ser73 of human JUN, which can detect the bands around 42-45 kDa.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



1X10^6 NIH/3T3 cells untreated (dashed lines) or treated with UV which intracellularly stained with 0.25 ug Coralite® Plus 647 Phospho-JUN (Ser73) Recombinant Antibody (CL647-80086, Clone:4A18) (red), or 0.25 ug Coralite® Plus 647 Rabbit IgG Isotype Control RecAb (CL647-98136, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.