

For Research Use Only

CoraLite® Plus 647-conjugated Phospho-P53 (Ser15) Recombinant antibody

Catalog Number: CL647-80195



Basic Information

Catalog Number: CL647-80195	GenBank Accession Number: BC003596	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 7157	CloneNo.: 2J21
Source: Rabbit	UNIPROT ID: P04637	Recommended Dilutions: IF/ICC 1:50-1:500
Isotype: IgG	Full Name: tumor protein p53	Excitation/Emission maxima wavelengths: 654 nm / 674 nm
	Calculated MW: 44 kDa	
	Observed MW: 53 kDa	

Applications

Tested Applications: IF/ICC, FC (Intra)	Positive Controls: IF/ICC : etoposide treated HT-29 cells,
Species Specificity: human	

Background Information

P53 is a 53 kDa protein that is activated in response to alteration of normal cell homeostasis, including DNA damage, nutrient starvation, heat shock, virus infection, pH change, hypoxia, and oncogene activation. P53 maintains genetic stability by regulating different processes, such as cell-cycle arrest, DNA synthesis and repair, programmed cell death, and energy metabolism. In non-stressed conditions these proteins bind p53, ubiquitylate it and target it for degradation by the proteasome. In stressed conditions the function of the Mdm2-Mdm4 complex is blocked by phosphorylation, protein-binding events and/or enhanced degradation. (PMID: 19935675, PMID: 24379683)

Storage

Storage:
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ A431 cells untreated (dashed line) or UV treated A431 cells (red) were intracellularly stained with 0.5 ug CoraLite® Plus 647 Anti-Human Phospho-P53 (Ser15) (CL647-80195, Clone:2J21), or 0.5 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.



Immunofluorescent analysis of (4% PFA) fixed etoposide treated HT-29 cells using CoraLite® Plus 647 Phospho-P53 (Ser15) antibody (CL647-80195, Clone: 2J21) at dilution of 1:200.