For Research Use Only

## CoraLite® Plus 647 Anti-Human Granzyme B Rabbit Recombinant Antibody



**Purification Method:** 

Protein A purification

Excitation/Emission maxima

CloneNo.:

240761G4

wavelengths: 654 nm / 674 nm

Catalog Number: CL647-98070

**Basic Information** 

Catalog Number: CL647-98070

100tests, 5 ul/test

Source: Rabbit

Isotype:

GenBank Accession Number:

GeneID (NCBI):

3002

**UNIPROT ID:** P10144 Full Name:

BC030195

granzyme B (granzyme 2, cytotoxic T-

lymphocyte-associated serine esterase 1)

Calculated MW: 247 aa, 28 kDa

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity:

human

## **Background Information**

GZMB (Granzyme B) is also named as CGL1, CSPB, CTLA1, GRB and belongs to the Granzyme subfamily. This enzyme is necessary for target cell lysis in cell-mediated immune responses. The cytotoxic lymphocyte protease granzyme B (GzmB) can promote apoptosis through direct processing and activation of members of the caspase family. GzmB can also cleave the BH3-only protein, BID, to promote caspase-independent mitochondrial permeabilization (PMID:17283187). GzmB induces lamin B degradation in isolated nuclei less efficiently than GzmA (PMID:11331782). This full-length protein has 2 glycosylation sites and a signal peptide. Unglycosylated human granzyme B is 26 kDa and high mannose glycosylated is 32 kDa and only 32 kDa or smaller forms of granzyme B are accumulated within nuclei (PMID:8626751). GzmB also forms dimers.


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 0.09% sodium azide and 0.5% BSA

in USA), or 1(312) 455-8498 (outside USA)

## **Selected Validation Data**



1x10^6 human PBMCs were intracellularly stained with FITC Plus Anti-Human CD8a, and 5 ul CoraLite® Plus 647 Anti-Human Granzyme B Rabbit RecAb (CL647-98070, Clone: 240761G4) or CoraLite® Plus 647 Rabbit IgG Isotype Control RecAb (CL647-98136, Clone: 240953C9). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



CL647-98070(Granzyme B),APC-A

1x10^6 human PBMCs were intracellularly stained with 5 ul Coralite® Plus 647 Anti-Human Granzyme B Rabbit RecAb (CL647-98070, Clone: 240761G4) (red), or Coralite® Plus 647 Rabbit IgG Isotype Control RecAb (CL647-98136, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).