For Research Use Only

CoraLite® Plus 647 Anti-Rat IL-4 Rabbit Recombinant Antibody

Catalog Number: CL647-98255

Basic Information	Catalog Number: CL647-98255	GenBank Accession Number: NM_201270.1	Purification Method: Protein A purification
	Size: 100ug , 500 µg/ml	GenelD (NCBI): 287287	CloneNo.: 241249H7
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: EG1113	UNIPROT ID: P20096 Full Name: interleukin 4	Recommended Dilutions: FC (Intra): 0.25 ug per 10^6 cells in a 100 µl suspension Excitation/Emission maxima
		Applications	Tested Applications: FC (Intra)
Species Specificity: rat			
Background Information	IL-4 is a cytokine produced by CD4+ T cells in response to helminthes and other extracellular parasites. It promotes the proliferation and differentiation of antigen presenting cells. IL-4 also plays a pivotal role in antibody isotype switching and stimulates the production of IgE. This cytokine has been applied in the treatment of autoimmune disorder like multiple myeloma, cancer, psoriasis, and arthritis. IL-4 has also been extensively applied to inhibit detrimental effect of Th1. It may promote the growth of epithelial tumors by mediating increased proliferation and survival.		
Storage	Storage: Store at 2-8°C. Avoid exposure to Storage Buffer: PBS with 0.09% sodium azide, pH	light. Stable for one year after shipme 7.3	ent.

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

www.ptglab.com

Selected Validation Data



1x10^{^6} untransfected or IL-4 transfected HEK-293T cells were intracellularly stained with 0.25 ug CoraLite® Plus 647 Anti-Rat IL-4 Rabbit RecAb (CL647-98255, Clone: 241249H7). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). 1x10^6 untransfected or IL-4 transfected HEK-293T cells were intracellularly stained with 0.25 ug CoraLite® Plus 647 Anti-Rat IL-4 Rabbit RecAb (CL647-98255, Clone: 241249H7) (red) or 0.25 ug CoraLite® Plus 647 Rabbit IgG Isotype Control RecAb (CL647-98136, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).