For Research Use Only

FcZero-rAb™ CoraLite® Plus 647 Anti-Mouse IL-17A Rabbit Recombinant Antibody



Catalog Number: CL647-FcA98005

Basic Information

Catalog Number: GenBank Accession Number: CL647-FcA98005 NM-010552

CL647-FcA98005 NM-010552 Protein A purification
Size: GeneID (NCBI): CloneNo.:

100ug, 500 ug/ml 16171 230038C7

Source: UNIPROT ID: Recommended Dilutions:

Rabbit Q62386 FC (Intra): 0.25 ug per 10^6 cells in a

Isotype: Full Name: 100 µl suspension

gG interleukin 17A Excitation/Emission maxima
mmunogon Catalog Number Calculated MW: wavelengths:

Immunogen Catalog Number: Calculated MW: wavelengths: EG0624 17 kDa 654 nm / 674 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity:

mouse

Positive Controls:

FC (Intra): PMA and ionomycin treated BALB/C Th17-

Purification Method:

polarized BALB/C splenocytes,

Background Information

IL-17A, also named as IL-17, is a proinflammatory cytokine. IL-17, synthesized only by memory T cells and natural killer cells, has pleiotropic effects, mainly in the recruitment and activation of neutrophils. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. The IL-17 receptor is a type I transmembrane protein, that is widely expressed on epithelial cells, fibroblasts, B and T cells, and monocytic cells. In psoriatic skin lesions, both Th17 cells and their downstream effector molecules, e.g. IL-17 and IL-22, are highly increased.

Storage


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer

PBS with 0.09% sodium azide, pH7.3

Selected Validation Data



1x10^6 untreated or PMA and ionomycin treated BALB/C Th17-polarized BALB/C splenocytes were intracellularly stained with PE Anti-Mouse CD4 and 0.25 ug CoraLite® Plus 647 Anti-Mouse IL-17A Rabbit RecAb (CL647-FcA98005, Clone:230038C7). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.