

For Research Use Only

CoraLite®650-conjugated c-MYC Polyclonal antibody



Catalog Number: CL650-10828

Featured Product

Basic Information

Catalog Number: CL650-10828	GenBank Accession Number: BC000141	Purification Method: Antigen affinity purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 4609	Excitation/Emission maxima wavelengths: 654 nm / 674 nm
Source: Rabbit	Full Name: v-myc myelocytomatosis viral oncogene homolog (avian)	
Isotype: IgG	Calculated MW: 49 kDa	
Immunogen Catalog Number: AG1263	Observed MW: 62-65 kDa, 50 kDa	

Applications

Tested Applications:
FC (Intra)

Species Specificity:
human

Background Information

MYC contains one basic helix-loop-helix (bHLH) domain. This protein is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. It seems to activate the transcription of growth-related genes. It binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt Lymphoma. This antibody is a rabbit polyclonal antibody raised against recombinant protein of human MYC. The 50kDa band recognized by antibody is the native form of MYC, while the other bands, between 60-70kDa, are the phosphorylated form of MYC [PMID: 12189186]

Storage

Storage:
Store at -20°C. Avoid exposure to light.


Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1×10^6 HeLa cells were intracellularly stained with 0.2 μ g CoraLite®650 Anti-Human c-MYC (CL650-10828) (red), or 0.2 μ g Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).