For Research Use Only

CoraLite®700 Anti-Human CD8 (UCHT4) Mouse IgG2a Recombinant Antibody




Catalog Number:CL700-65559


Basic Information	Catalog Number: CL700-65559	GenBank Accession Number: BC025715	Purification Method: Protein A purification					
	Size: 100tests , 5 ul/test	GenelD (NCBI): 925	CloneNo.: UCHT4					
	Source: Mouse Isotype: IgG2a	ENSEMBL Gene ID: ENSG00000153563 UNIPROT ID: P01732 Full Name: CD8a molecule Calculated MW: 235 aa, 26 kDa	Recommended Dilutions: FC: 5 ul per 10^6 cells in a 100 µl suspension Excitation/Emission maxima wavelengths: 702 nm / 723 nm					
				Applications	Tested Applications:	Positive C	Positive Controls:	
					FC Species Specificity: human			
				Background Informatior	CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer o CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8aβ heterodimer while CD8aa homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) γδ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).			
	been found on some gut intra 2111591; 1831127; 8420975) activation (PMID: 8499079). [epithelial T cells , on some T cell receptor . CD8 acts as a co-receptor that binds to MH	(TCR) $\gamma \delta$ T cells and on NK cells (PMID: C class-I and participates in cytotoxic T cell					

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10^6 human PBMCs were surface stained with CoraLite® Plus 488 Anti-Human CD3, and 5 ul CoraLite® 700 Anti-Human CD8 (UCHT4) Mouse IgG2a RecAb (CL700-65559, Clone: UCHT4) or unstained. Cells were not fixed.

1x10[^]6 human PBMCs were surface stained with 5 ul CoraLite®700 Anti-Human CD8 (UCHT4) Mouse IgG2a RecAb (CL700-65559, Clone: UCHT4) (red) or unstained (blue). Cells were not fixed.