For Research Use Only

CoraLite® Plus 750-conjugated PCNA Monoclonal antibody



Purification Method:

CloneNo.:

10D10E11

Protein G purification

Recommended Dilutions:

Excitation/Emission maxima

WB 1:2000-1:16000

wavelengths: 755 nm / 780 nm

Catalog Number: CL750-60097

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL750-60097 BC000491 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 5111 Nanodrop: **UNIPROT ID:** P12004

Mouse Full Name:

Isotype: proliferating cell nuclear antigen lgG1

Calculated MW: Immunogen Catalog Number: 29 kDa/31 kDa AG7416 Observed MW:

36-38 kDa

Applications

Tested Applications:

Species Specificity: human, mouse, rat, pig Positive Controls:

WB: HeLa cells, HepG2 cells

Background Information

Proliferating Cell Nuclear Antigen, commonly known as PCNA, is a protein that acts as a processivity factor for DNA polymerase δ in eukaryotic cells. This protein is an auxiliary protein of DNA polymerase delta and is involved in the control of eukaryotic DNA replication by increasing the polymerase's processibility during elongation of the leading strand. PCNA induces a robust stimulatory effect on the 3'-5' exonuclease and 3'-phosphodiesterase, but not apurinic-apyrimidinic (AP) endonuclease, APEX2 activities. It has to be loaded onto DNA in order to be able to stimulate APEX2. PCNA protein is highly conserved during evolution; the deduced amino acid sequences of rat and human differ by only 4 of 261 amino acids. PCNA has been used as loading control for proliferating cells. The calculated molecular weight of PCNA is 29 kDa, but modified PCNA is 36kDa (PMID: 1358458).

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with CL750-60097 (PCNA antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.