

For Research Use Only

CoraLite® Plus 750-conjugated CD99 Monoclonal antibody

Catalog Number: CL750-60354



Basic Information

Catalog Number:

CL750-60354

Size:

100ul , Concentration: 1000 ug/ml by
Nanodrop;

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG19474

GenBank Accession Number:

BC021620

GeneID (NCBI):

4267

UNIPROT ID:

P14209

Full Name:

CD99 molecule

Calculated MW:

19 kDa, 16 kDa, 17 kDa

Observed MW:

32 kDa

Purification Method:

Protein G purification

CloneNo.:

5B6A4

Recommended Dilutions:

WB 1:500-1:1000

Excitation/Emission maxima
wavelengths:

755 nm / 780 nm

Applications

Tested Applications:

WB

Species Specificity:

human

Positive Controls:

WB : Raji cells,

Background Information

CD99, also known as MIC2, is a heavily O-glycosylated transmembrane protein involved in T-cell adhesion processes and in spontaneous rosette formation with erythrocytes. CD99 is broadly distributed on many cell types, with particularly strong expression on human cortical thymocytes, Ewing's sarcoma cells and peripheral primitive neuroectodermal tumors (PMID: 9794396; 16984917). In normal cells, CD99 has been functionally implicated in cell adhesion, migration, apoptosis, differentiation, activation, and proliferation of lymphocytes and monocyte extravasation and transport of several transmembrane proteins (PMID: 16984917). CD99 displays two surface isoforms generated by alternative splicing: a long 32 kDa and a short 28 kDa form (PMID: 12368226).

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with CL750-60354 (CD99 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.