For Research Use Only

CoraLite® Plus 750 Anti-Human CD3 (OKT3)



Purification Method:

Affinity purification

Excitation/Emission maxima

CloneNo.:

wavelengths:

755 nm / 780 nm

OKT3

Catalog Number: CL750-65133

Basic Information

Catalog Number: CL750-65133

BC049847

GeneID (NCBI): 100tests, 5 µl/test 916

Full Name:

CD3e molecule, epsilon (CD3-TCR Mouse complex) Isotype: Calculated MW: IgG2a, kappa

207 aa, 23 kDa

GenBank Accession Number:

Applications

Tested Applications:

Species Specificity:

Background Information

CD3 is a multimeric protein associated with the T-cell receptor (TCR) to form a complex involved in antigen recognition and signal transduction (PMID: 15885124). CD3 is composed of CD3 γ , δ , ϵ , and ζ chains (PMID: 1826255). It is expressed by thymocytes in a developmentally regulated manner, T cells, and some NK cells (PMID: 3289580). The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction (PMID: 11985657).

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 0.09% sodium azide and 0.5% BSA.

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^6 human PBMCs were surface co-stained with FITC Anti-Human CD19 and 5 ul CoraLite® Plus 750 Anti-Human CD3 (CL750-65133, Clone:OKT3) or Mouse IgG2a Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul Coralite® Plus 750 Anti-Human CD3 (CL750-65133, Clone:OKT3) or Mouse IgG2a Isotype Control. Cells were not fixed. Lymphocytes were gated.