For Research Use Only

CoraLite® Plus 750-conjugated Lamin B1 Monoclonal antibody

Catalog Number:CL750-66095

Featured Product




| Basic Information | Catalog Number: CL750-66095 | GenBank Accession Number: BC012295 | Purification Method: Protein G purification |
|---|---|---|---|
| | Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG20522 | GeneID (NCBI): 4001 | CloneNo.: 3C10G12 |
| | | ENSEMBL Gene ID: ENSG00000113368 | Recommended Dilutions: WB 1:2000-1:16000 |
| | | UNIPROT ID: P20700 Full Name: lamin B1 | Excitation/Emission maxima wavelengths: 755 nm / 780 nm |
| | | | |
| | | Observed MW: 66-70 kDa | |
| | | Applications | |
| WB Species Specificity: human, mouse, rat | WB : HeL | | a cells, HepG2 cells |
| Background Information | Lamins are components of the nuclear lamina, a fibrous layer on the nucleoplasmic side of the inner nuclear membrane, which is thought to provide a framework for the nuclear envelope and may also interact with chromatin. The nuclear lamina consists of a two-dimensional matrix of proteins located next to the inner nuclear membrane. The lamin family of proteins make up the matrix and are highly conserved in evolution. During mitosis, the lamina matrix is reversibly disassembled as the lamin proteins are phosphorylated. Vertebrate lamins consist of two types, A and B. This gene encodes one of the two B type proteins, B1. This protein is not suitable for samples where the nuclear envelope has been removed. | | |
| Storage | Storage: Store at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% Glycerol, 0.05% Procli Aliquoting is unnecessary for -20°C s | n300, 0.5% BSA, pH 7.3. | ent. |

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with CL750-66095 (Lamin B1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.