For Research Use Only

## CoraLite® Plus 750 Anti-Mouse ICAM-1/CD54 Rabbit Recombinant Antibody

Catalog Number: CL750-98049



**Basic Information** 

Catalog Number:

NM\_010493

**Purification Method:** Protein A purification

CL750-98049 Size:

GeneID (NCBI):

GenBank Accession Number:

CloneNo.:

100ug , 500  $\mu$ g/ml

15894

240249D10

Source: Rabbit

**UNIPROT ID:** P13597

Excitation/Emission maxima

Isotype:

Full Name:

wavelengths: 755 nm / 780 nm

intercellular adhesion molecule 1

EG0482

Calculated MW: Immunogen Catalog Number:

59 kDa

**Applications** 

**Tested Applications:** 

**Positive Controls:** 

Species Specificity:

mouse

FC: mouse splenocytes,

## **Background Information**

ICAM-1 (CD54) is a 90-kDa transmembrane glycoprotein of the immunoglobulin superfamily and is critical for the firm attachment and transmigration of leukocytes out of blood vessels and into tissues (PMID: 19307690). ICAM-1 is expressed by several cell types, typically on endothelial cells and cells of the immune system, and its expression can be up-regulated by various stimuli, including TNF-a, INF-y, IL-1 and thrombin (PMID: 3086451; 9694714; 15979056). It is a ligand for LFA-1 and Mac-1, serves as a receptor for rhinovirus, and is one of several receptors used by Plasmodium falciparum (PMID: 2566624; 2538244; 2475784).

Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3


in USA), or 1(312) 455-8498 (outside USA)

## Selected Validation Data





1x10^6 mouse splenocytes were surface stained with CoraLite® Plus 488 Anti-Mouse CD3, and 0.25 ug CoraLite® Plus 750 Anti-Mouse ICAM-1/CD54 Rabbit Recombinant Antibody (CL750-98049, Clone: 240249D10) or Isotype Control. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 mouse splenocytes were surface stained with 0.25 ug CoraLite® Plus 750 Anti-Mouse ICAM-1/CD54 Rabbit Recombinant Antibody (CL750-98049, Clone: 240249D10) (red) or 0.25 ug Isotype Control (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.