For Research Use Only

FcZero-rAb™ CoraLite® Plus 750 Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2) Rabbit IgG Recombinant Antibody



Catalog Number: CL750-FcA65663

Basic Information

Catalog Number: GenBank Accession Number: **Purification Method**: CL750-FcA65663 BC031711 Protein A purification

Size: GeneID (NCBI): CloneNo: 100ug, 500 ug/ml M5/114.15.2

UNIPROT ID: Source: Recommended Dilutions:

Rabbit FC: 0.25 ug per 10^6 cells in a 100 µl P14434 suspension

Isotype: **Full Name:**

histocompatibility 2, class II antigen Excitation/Emission maxima IgG wavelengths:

A. alpha 755 nm / 780 nm

Applications

Tested Applications: Positive Controls:

FC: mouse splenocytes, Species Specificity:

mouse

Background Information

Major histocompatibility complex (MHC) class II molecules are heterodimeric transmembrane glycoproteins that are expressed on the surface of antigen-presenting cells such as dendritic cells, macrophages, and B cells. The M5/114.15.2 antibody detects polymorphic determinants present on B cells, monocytes, macrophages, dendritic cells, and activated T lymphocytes from mice carrying the H-2b, H-2d, H-2q, H-2p, H-2r and H-2u haplotypes but not from mice carrying the H-2s or H-2f haplotypes (PMID: 6170707). This antibody recognizes both I-A and I-E subregion-encoded Ia glycoproteins (I-Ab, I-Ad, I-Ad, I-Ed, I-Ek, not I-Af, I-Ak, or I-As). M5/114 antibody is reported to inhibit I-A subregion-restricted T cell responses of the H-2b, H-2d, H-2q, H-2u but not the H-2f, H-2k, H-2s haplotypes (PMID: 6173436).


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer

PBS with 0.09% sodium azide, pH7.3

Selected Validation Data



1x10^6 mouse splenocytes were surface stained with PE Anti-Mouse CD3e (145-2C11), and 0.25 ug Coralite® Plus 750 Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2) Rabbit IgG RecAb (CL750-FcA65663, Clone: M5/114.15.2) or 0.25 ug Rabbit IgG Isotype Control RecAb (CL750-FcA98136, Clone: 240953C9). Cells were not fixed.



CL750-FcA65663(MHC Class II (I-A/I-E)),APC-A750-A

1x10^6 mouse splenocytes were surface stained with 0.25 ug CoraLite® Plus 750 Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2) Rabbit IgG RecAb (CL750-FcA65663, Clone: M5/114.15.2) (red) or 0.25 ug Rabbit IgG Isotype Control RecAb (CL750-FcA98136, Clone: 240953C9) (blue). Cells were not fixed