For Research Use Only

FcZero-rAb® CoraLite® Plus 750 Anti-Mouse CD8a Rabbit Recombinant Antibody

Catalog Number:CL750-FcA98472-2

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Basic Information

Catalog Number:
CL750-FcA98472-2

Size:
100ug , 500 ug/ml

Source:
Rabbit

Isotype:
IgG

Immunogen Catalog Number:
EG3265

GenBank Accession Number:
NM_001081110.2

GeneID (NCBI):
12525

UNIPROT ID:
P01731

Full Name:
CD8 antigen, alpha chain

Calculated MW:
27 kDa

Purification Method:
Protein A purification

CloneNo.:
242879C10

Recommended Dilutions:
FC: 0.25 ug per 10^6 cells in a 100 µl suspension

Excitation/Emission maxima wavelengths:
755 nm / 780 nm

Applications

Tested Applications:
FC

Species Specificity:
mouse

Positive Controls:

FC : mouse splenocytes,

Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8α or as a heterodimer of CD8α and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8αβ heterodimer while CD8αα homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) γδ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).


Storage

Storage:
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 0.09% sodium azide, pH7.3


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10^6 mouse splenocytes were surface stained with FcZero-rAb® CoraLite® Plus 488 Anti-Mouse CD3, and 0.25 ug CoraLite® Plus 750 Anti-Mouse CD8a Rabbit RecAb (CL750-FcA98472-2, Clone: 242879C10) or 0.25 ug FcZero-rAb® CoraLite® Plus 750 Rabbit IgG Isotype Control Recombinant Antibody (CL750-FcA98136, Clone: 240953C9). Cells were not fixed.



1x10^6 mouse splenocytes were surface stained with 0.25 ug CoraLite® Plus 750 Anti-Mouse CD8a Rabbit RecAb (CL750-FcA98472-2, Clone: 242879C10) (red) or 0.25 ug FcZero-rAb® CoraLite® Plus 750 Rabbit IgG Isotype Control Recombinant Antibody (CL750-FcA98136, Clone: 240953C9) (blue). Cells were not fixed.