

For Research Use Only

FcZero-rAb® PerCP-Cyanine5.5 Anti-Mouse CD25/IL-2RA Rabbit Recombinant Antibody

Catalog Number:CPY5-FcA98288



Basic Information

Catalog Number:

CPY5-FcA98288

Size:

100ug, 200 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG3087

GenBank Accession Number:

NM_008367.3

GeneID (NCBI):

16184

UNIPROT ID:

P01590

Full Name:

interleukin 2 receptor, alpha chain

Calculated MW:

31 kDa

Purification Method:

Protein A purification

CloneNo.:

242494H5

Recommended Dilutions:

FC: 0.25 ug per 10⁶ cells in a 100 µl suspension

Excitation/Emission maxima wavelengths:

482 nm / 690nm

Applications

Tested Applications:

FC

Species Specificity:

mouse

Positive Controls:

FC : anti-CD3/CD28 treated mouse splenocytes,

Background Information

Proliferation of T lymphocytes is triggered by the interaction of IL-2 with its specific receptor following T lymphocyte activation. The receptor for IL-2 has three forms, generated by different combinations of three different proteins, the alpha chain (IL-2R alpha), the beta chain (IL-2R beta), and the gamma chain (IL-2R gamma) (PMID: 8476561). IL-2R alpha (also known as CD25) is a type I transmembrane protein present on activated T cells, activated B cells, some thymocytes, myeloid precursors, and oligodendrocytes. CD25 is also found on natural CD4⁺Foxp3⁺ Treg cells and a subset of human CD4⁺ memory T cells (PMID: 22585674).

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ untreated or anti-CD3/CD28 treated mouse splenocytes were surface stained with FcZero-rAb® CoraLite® Plus 488 Anti-Mouse CD4 and 0.25 ug PerCP-Cyanine5.5 Anti-Mouse CD25/IL-2RA Rabbit RecAb (CPY5-FcA98288, Clone: 242494H5). Cells were incubated with MonoZero™ Monocytes blocking Reagent (PF00020) prior to staining. Cells were not fixed.

1x10⁶ anti-CD3/CD28 treated mouse splenocytes were surface stained with 0.25 ug PerCP-Cyanine5.5 Anti-Mouse CD25/IL-2RA Rabbit RecAb (CPY5-FcA98288, Clone: 242494H5) (red) or 0.25 ug FcZero-rAb® PerCP-Cyanine5.5 Rabbit IgG Isotype Control Recombinant Antibody (CPY5-FcA98136, Clone: 240953C9) (blue). 1x10⁶ untreated mouse splenocytes were surface stained with 0.25 ug PerCP-Cyanine5.5 Anti-Mouse CD25/IL-2RA Rabbit RecAb (CPY5-FcA98288,

