

For Research Use Only

# Recombinant Human BAFF/TNFSF13B protein (hFc Tag)

Catalog Number: Eg0155



## Basic Information

**Species:**  
Human

**EC50:**  
0.6-2.4 ng/mL

**Purity:**  
>90 %, SDS-PAGE

**Tag:**  
hFc Tag

## Technical Specifications

**Purity:**  
>90 %, SDS-PAGE

**Endotoxin Level:**  
<0.1 EU/μg protein, LAL method

**Source:**  
HEK293-derived Human BAFF protein Ala134-Leu285 (Accession# Q9Y275-1) with a human IgG1 Fc tag at the N-terminus.

**GeneID:**  
10673

**Accession:**  
Q9Y275-1

**Predicted Molecular Mass:**  
43 kDa

**SDS-PAGE:**  
45-50 kDa, reducing (R) conditions

**Formulation:**  
Lyophilized from 0.22 μm filtered solution in PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.

## Biological Activity

Immobilized Human BCMA (Myc tag, His tag) at 2 μg/mL (100 μL/well) can bind Human BAFF (hFc tag) with a linear range of 0.6-2.4 ng/mL.

## Storage and Shipping

**Storage:**  
It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

- Until expiry date, -20°C to -80°C as lyophilized proteins.
- 3 months, -20°C to -80°C under sterile conditions after reconstitution.

**Shipping:**  
The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.

## Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

## Background

TNFSF13B, also known as BAFF, BLYS, TALL1, TNFSF20, ZTNF4 and CD257, is a cytokine that binds to TNFRSF13B/TACI and TNFRSF17/BCMA. This cytokine is expressed in B lineage cells, and acts as a potent B cell activator. The BAFF system promotes B cell survival and differentiation and plays a prominent role in the pathogenesis of autoimmune diseases. It has been shown that BAFF inhibitors are in clinical trials for systemic lupus erythematosus with significant efficacy.

## References

1. Cristian R Smulski et al (2018). Front Immunol. Oct 8;9:2285.
2. Edina Schweighoffer et al (2021). Curr Opin Immunol. Aug;71:124-131.
3. Anne Davidson. (2010). Curr Opin Immunol. Dec;22(6):732-9.
4. Shingo Nakayamada et al (2016). Inflamm Regen. Jul 21;36:6.

## Synonyms


TNFSF13B, B cell activating factor, B lymphocyte stimulator, BAFF, B-cell-activating factor

For technical support and original validation data for this product please contact


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Purity of Recombinant Human BAFF was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) and non-reducing (NR) conditions and stained using Coomassie blue.



Immobilized Human BCMA (Myc tag, His tag) at 2 µg/mL (100 µL/well) can bind Human BAFF (hFc tag) with a linear range of 0.6-2.4 ng/mL