For Research Use Only

# Recombinant Human Siglec-7/CD328 protein (His Tag)

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Catalog Number: Eg1545

## Basic Information

**Species:**
Human

**Purity:**
>90 %, SDS-PAGE

**Tag:**
His Tag

## Technical Specifications

**Purity:**
>90 %, SDS-PAGE

**Endotoxin Level:**
<1.0 EU/μg protein, LAL method

**Source:**
HEK293-derived Human Siglec-7/CD328 protein Gln19-Leu353 (Accession# Q9Y286-1) with a His tag at the C-terminus.

**GeneID:**
27036

**Accession:**
Q9Y286-1

**Predicted Molecular Mass:**
38.0 kDa

**SDS-PAGE:**

**Formulation:**
Lyophilized from sterile PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.

## Biological Activity

Not tested

## Storage and Shipping

**Storage:**
It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

- Until expiry date, -20°C to -80°C as lyophilized proteins.
- 3 months, -20°C to -80°C under sterile conditions after reconstitution.

**Shipping:**
The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.

## Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

## Background

Sialic acid binding Ig-like lectin 7 (Siglec-7), also known as CD328 or p75/AIRM-1, is a member of the Siglec family of glycan-recognition proteins. Siglec-7 is a type-I transmembrane protein consisting of three extracellular immunoglobulin-like domains that comprise an N-terminal V-set domain and two C2-set domains, a transmembrane region and a cytoplasmic tail containing two tyrosine residues embodied in immunoreceptor tyrosine-based inhibition motif-like motifs. It is mainly expressed on immune cells, with low levels on granulocytes, intermediate levels on monocytes, and relatively high levels on a major subset of natural killer cells and a minor subset of CD8+ T cells. Siglec-7 is an inhibitory receptor that negatively regulates the function of NK cells and modulates the immune response through the interaction of sialic acid-containing ligands .

## References


1. Zheng, Yayun et al. Journal of immunology research vol. 2020 6243819.
2. Nicoll, G et al. The Journal of biological chemistry vol. 274,48 (1999): 34089-95.
3. Shao, J-Y et al. Scandinavian journal of immunology vol. 84,3 (2016): 182-90.


## Synonyms


**For technical support and original validation data for this product please contact**

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

## Selected Validation Data