

Recombinant Mouse CD38 protein (rFc Tag)


Catalog Number: Eg2915

Basic Information	Species: Mouse	Purity: >90 %, SDS-PAGE	Tag: rFc Tag
Technical Specifications	<p>Purity: >90 %, SDS-PAGE</p> <p>Endotoxin Level: <0.1 EU/μg protein, LAL method</p> <p>Source: HEK293-derived Mouse CD38 protein Leu45-Thr304 (Accession# P56528) with a rabbit IgG Fc tag at the C-terminus.</p> <p>GenID: 12494</p> <p>Accession: P56528</p> <p>Predicted Molecular Mass: 55.8 kDa</p> <p>SDS-PAGE: 60-72 kDa, reducing (R) conditions</p> <p>Formulation: Lyophilized from 0.22 μm filtered solution in PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.</p>		
Biological Activity	Not tested		
Storage and Shipping	<p>Storage: It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> Until expiry date, -20°C to -80°C as lyophilized proteins. 3 months, -20°C to -80°C under sterile conditions after reconstitution. <p>Shipping: The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.</p>		
Reconstitution	Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.		
Background	CD38, also known as ADP-ribosyl cyclase 1, is a type II transmembrane glycoprotein with a short N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites (PMID: 2319135). The extracellular domain of CD38 has bifunctional enzyme activities that catalyze synthesis of cyclic ADP ribose from nicotinamide adenine dinucleotide (NAD) and hydrolysis of cyclic ADP ribose to adenosine diphosphoribose (PMID: 10636863). CD38 is expressed on a variety of hematopoietic and non-hematopoietic cells and is involved in diverse processes such as generation of calcium-mobilizing metabolites, cell activation, and chemotaxis (PMID: 25938500).		
References	<ol style="list-style-type: none"> Jackson DG, Bell JI. (1990) J Immunol. 144(7):2811-5. Cho YS, Han MK, et al. (2000) J Biol Chem. 275(3):1685-90. Schneider M, Schumacher V, et al. (2015) PLoS One. 10(5):e0126007. 		
Synonyms	Cd38, CD38 antigen, Cd38 rs1		

For technical support and original validation data for this product please contact

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.comThis product is exclusively available under
Proteintech Group brand and is not available
to purchase from any other manufacturer.

Selected Validation Data



Purity of Recombinant Mouse CD38 was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) and non-reducing (NR) conditions and stained using Coomassie blue.