

For Research Use Only

FITC Plus Anti-Human IL-10 (7B8G3)

Catalog Number: FITC-65286



Basic Information

Catalog Number: FITC-65286	GenBank Accession Number: BC104252	Purification Method: Affinity purification
Size: 100ug, 0.2 mg/mL	GeneID (NCBI): 3586	CloneNo.: 7B8G3
Source: Mouse	Full Name: interleukin 10	Excitation/Emission maxima wavelengths: 495 nm / 524 nm
Isotype: IgG2b	Calculated MW: 21 kDa	
Immunogen Catalog Number: HZ-1145		

Applications

Tested Applications:
FC

Species Specificity:
Human

Background Information

Interleukin (IL)-10 is an anti-inflammatory cytokine, produced by T helper (Th) cells, macrophages, monocytes, and B cells, that plays a crucial role in preventing inflammatory and autoimmune pathologies. It downregulates the expression of Th1 cytokines, MHC class II antigens, and co-stimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. IL-10 can block NF- κ B activity, and is involved in the regulation of the JAK-STAT signaling pathway. IL-10, along with its receptors, describes an important role in pathogenesis of various diseases, including infectious, inflammatory, autoimmune diseases. IL-10 mutations are associated with an increased susceptibility to HIV-1 infection and rheumatoid arthritis.

Storage


Storage:
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:
PBS with 0.09% sodium azide and 0.5% BSA.


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ LPS and Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug FITC Plus Anti-Human IL-10 (FITC-65286, Clone:7B8G3) or Mouse IgG2b Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). Monocytes were gated.



1X10⁶ LPS and Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug FITC Plus Anti-Human IL-10 (FITC-65286, Clone:7B8G3) (red) or Mouse IgG2b Isotype Control (blue), or unactivated and Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug FITC Plus Anti-Human IL-10 (FITC-65286, Clone:7B8G3) (black, dashed). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit

