For Research Use Only

FITC Plus Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2) Rat IgG2a Recombinant Antibody



Catalog Number: FITC-65663

Catalog Number: **Basic Information**

FITC-65663

100ug , 500 ug/ml

Source:

Isotype:

lgG2a

GenBank Accession Number:

BC031711 GeneID (NCBI):

14960

UNIPROT ID: P14434

Full Name:

histocompatibility 2, class II antigen Excitation/Emission maxima

A, alpha

Purification Method:

Protein G purification

CloneNo.: M5/114.15.2

Recommended Dilutions:

FC: 0.25 ug per 10^6 cells in a 100 μ l

suspension

wavelengths: 495 nm / 524 nm

Applications

Tested Applications:

Species Specificity:

mouse

Positive Controls:

FC: mouse splenocytes,

Background Information

Major histocompatibility complex (MHC) class II molecules are heterodimeric transmembrane glycoproteins that are expressed on the surface of antigen-presenting cells such as dendritic cells, macrophages, and B cells. The M5/114.15.2 antibody detects polymorphic determinants present on B cells, monocytes, macrophages, dendritic cells, and activated T lymphocytes from mice carrying the H-2b, H-2d, H-2q, H-2p, H-2r and H-2u haplotypes but not from mice carrying the H-2s or H-2f haplotypes (PMID: 6170707). This antibody recognizes both I-A and I-E subregion-encoded Ia glycoproteins (I-Ab, I-Ad, I-Ad, I-Ed, I-Ek, not I-Af, I-Ak, or I-As). M5/114 antibody is reported to inhibit I-A subregion-restricted T cell responses of the H-2b, H-2d, H-2q, H-2u but not the H-2f, H-2k, H-2s haplotypes (PMID: 6173436).


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide, pH7.3

Selected Validation Data



1x10^6 mouse splenocytes were surface stained with APC Anti-Mouse CD3, and 0.25 ug FITC Plus Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2) Rat IgG2a RecAb (FITC-65663, Clone:M5/114.15.2) or FITC Plus Rat IgG2a Isotype Control (2A3) (FITC-65209, Clone: 2A3). Cells were not fixed.



FITC-65663 MHC Class II (I-A/I-E),FITC-A

1x10^6 mouse splenocytes were surface stained with 0.25 ug FITC Plus Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2) Rat IgG2a RecAb (FITC-65663, Clone: M5/114.15.2) (red) or FITC Plus Rat IgG2a Isotype Control (2A3) (FITC-65209, Clone: 2A3) (blue). Cells were not fixed.