For Research Use Only

FITC Plus Anti-Mouse IL-1 alpha Rabbit Recombinant Antibody

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Purification Method:

Protein A purification

Catalog Number: FITC-98052

Basic Information

Catalog Number: GenBank Accession Number: FITC-98052 NM_010554

 Size:
 GeneID (NCBI):
 CloneNo.:

 100ug, 500 ug/ml
 16175
 240431G10

Source: UNIPROT ID: Recommended Dilutions:

Rabbit P01582 FC (Intra): 0.25 ug per 10^6 cells in a

 Isotype:
 Full Name:
 100 μl suspension

 IgG
 interleukin 1 alpha
 Excitation/Emission maxima

Immunogen Catalog Number: Calculated MW: wavelengths:
EG0263 31kd 495 nm / 524 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity:

mouse

Positive Controls:

FC (Intra): LPS treated RAW 264.7 cells,

Background Information

IL-1 alpha is a dual function cytokine that can function both as a transcription factor and a damage-associated molecular pattern (DAMP) in innate immune responses. It is expressed by many cell types and its expression is increased when cells are exposed to growth factors, pro-inflammatory, and stress-related signals. In its pro-form, IL-1 alpha contains a nuclear localization signal that allows it to translocate to the nucleus and bind to DNA, resulting in either the increase or prohibition of pro-inflammatory cytokine expression. When cells are exposed to necrotic signals, mature IL-1 alpha is released from the cell lysates and triggers a strong pro-inflammatory response upon binding to its receptor on nearby cells. Due to its strongly pro-inflammatory nature, IL-1 alpha has been linked to tumor metastasis and lower median survival in several cancers (PMID: 27434011, 29247995).

Storage


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide, pH7.3

Selected Validation Data



1x10^6 untreated or LPS-treated RAW 264.7 cells were intracellularly stained with 0.25 ug FITC Plus Anti-Mouse IL-1 alpha Rabbit RecAb (FITC-98052, Clone: 240431G10). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set.



1x10^6 LPS-treated RAW 264.7 cells were intracellularly stained with 0.25 ug FITC Plus Anti-Mouse IL-1 alpha Rabbit RecAb (FITC-98052, Clone: 240431G10) (red) or 0.25 ug FITC Plus Rabbit 1gG Isotype Control RecAb (FITC-98136, Clone: 240953C9) (blue). 1x10^6 untreated RAW 264.7 cells were intracellularly stained with 0.25 ug FITC Plus Anti-Mouse IL-1 alpha Rabbit RecAb (FITC-98052, Clone: 240431G10) (black, dashed). Cells were fixed and permeabilized with