For Research Use Only

## FITC Plus Anti-Human TNF-alpha Rabbit Recombinant Antibody

Catalog Number: FITC-98064



**Basic Information** 

Catalog Number:

FITC-98064

Size:

100tests, 5 ul/test

Source: Rabbit

Isotype:

GenBank Accession Number:

NM\_000594.4 GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000232810 **UNIPROT ID:** 

P01375

Full Name:

tumor necrosis factor (TNF superfamily, member 2)

Calculated MW:

26 kDa

**Purification Method:** 

Protein A purification

CloneNo.: 240668B11

Excitation/Emission maxima

wavelengths: 495 nm / 524 nm

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity:

**Background Information** 


TNF, as also known as TNF-alpha, or cachectin, is a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. It is expressed as a 26 kDa membrane bound protein and is then cleaved by TNF-alpha converting enzyme (TACE) to release the soluble 17 kDa monomer, which forms homotrimers in circulation. It is produced chiefly by activated macrophages, although it can be produced by many other cell types such as CD4+ lymphocytes, NK cells, neutrophils, mast cells, eosinophils, and neurons. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, ins resistance, and cancer.

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA

## **Selected Validation Data**



1x10^6 untreated or PMA, Ionomycin and Brefeldin A treated human PBMCs were intracellularly stained with CoraLite® Plus 647 Anti-Human CD4 and 5 ul FITC Plus Anti-Human TNF-alpha Rabbit RecAb (FITC-98064, Clone:240668B11). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



FITC-98064(TNF-alpha),FITC-

1x10^6 PMA, Ionomycin and Brefeldin A treated human PBMCs were intracellularly stained with 5 ul FITC Plus Anti-Human TNF-alpha Rabbit RecAb (FITC-98064, Clone: 240668B11) (red), or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9) (blue). 1x10^6 untreated human PBMCs were intracellularly stained with 5 ul FITC Plus Anti-Human TNF-alpha Rabbit RecAb (FITC-98064, Clone: 240668B11) (black, dashed). Cells were fixed with 4%



