For Research Use Only

## FITC Plus Anti-Human Granzyme B Rabbit Recombinant Antibody

Catalog Number: FITC-98070



**Purification Method:** 

CloneNo.:

240761G4

wavelengths:

495 nm / 524 nm

Protein A purification

Excitation/Emission maxima

**Basic Information** 

Catalog Number:

FITC-98070

Size:

100tests, 5 ul/test

Source: Rabbit

Isotype: IgG

GenBank Accession Number:

BC030195 GeneID (NCBI):

3002 **UNIPROT ID:** 

P10144 **Full Name:** 

granzyme B (granzyme 2, cytotoxic T-

lymphocyte-associated serine esterase 1)

Calculated MW: 247 aa, 28 kDa

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity:

human

## **Background Information**


GZMB (Granzyme B) is also named as CGL1, CSPB, CTLA1, GRB and belongs to the Granzyme subfamily. This enzyme is necessary for target cell lysis in cell-mediated immune responses. The cytotoxic lymphocyte protease granzyme  $B \ (GzmB) \ can promote \ apoptosis \ through \ direct \ processing \ and \ activation \ of \ members \ of \ the \ caspase \ family. \ GzmB$ can also cleave the BH3-only protein, BID, to promote caspase-independent mitochondrial permeabilization (PMID:17283187). GzmB induces lamin B degradation in isolated nuclei less efficiently than GzmA (PMID:11331782). This full-length protein has 2 glycosylation sites and a signal peptide. Unglycosylated human granzyme B is 26 kDa and high mannose glycosylated is 32 kDa and only 32 kDa or smaller forms of granzyme B are accumulated within nuclei (PMID:8626751). GzmB also forms dimers.

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA

## **Selected Validation Data**



1x10^6 human PBMCs were intracellularly stained with APC Anti-Human CD8a, and 5 ul FITC Plus Anti-Human Granzyme B Rabbit RecAb (FITC-98070, Clone: 240761G4) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



1x10^6 human PBMCs were intracellularly stained with 5 ul FITC Plus Anti-Human Granzyme B Rabbit RecAb (FITC-98070, Clone: 240761G4) (red) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).