For Research Use Only

FITC Plus Anti-Mouse CD48 Rabbit Recombinant Antibody

Proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Purification Method:

Catalog Number: FITC-98081

Basic Information

Catalog Number: GenBank Accession Number:

FITC-98081 BAE96326.1 Protein A purification

 Size:
 GeneID (NCBI):
 CloneNo.:

 100ug , 500 μg/ml
 12506
 240732A5

Source: UNIPROT ID: Recommended Dilutions:

Rabbit P18181 FC: 0.25 ug per 10^6 cells in a 100 µl

 Isotype:
 Full Name:
 suspension

 IgG
 CD48 antigen
 Excitation/Emiss

gG CD48 antigen Excitation/Emission maxima mmunogen Catalog Number: Calculated MW: wavelengths:

Immunogen Catalog Number: Calculated MW: Wavelengths: EG1323 27 kDa 495 nm / 524 nm

Applications

Tested Applications:

Species Specificity:

mouse

Positive Controls:

FC: mouse splenocytes,

Background Information

Signaling lymphocytic activation molecule family 2 (SLAMF2, CD48) is an adhesion and costimulatory molecule expressed constitutively on most hematopoietic cells, particularly in antigen-presenting cells (APC). CD48 is expressed on all the hematopoietic cells, both in humans and mice, except for murine neutrophils and long-term HSC. CD48 exists as both a membrane-bound (mCD48) and a soluble (sCD48) form. CD48 can activate T cells, antigen-presenting cells and granulocytes, by binding to CD2 or bacterial FimH, and through cell intrinsic effects. Interactions between CD48 and its high-affinity ligand CD244 are more complex, with both stimulatory and inhibitory outcomes. CD48 expression is increased in autoimmunity and allergy diseases, and anti-CD48 monoclonal antibodies have been shown to attenuate experimental autoimmune encephalomyelitis.

Storage


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 0.09% sodium azide, pH7.3

Selected Validation Data





1x10^6 mouse splenocytes were surface stained with Coralite® Plus 647 Anti-Mouse CD19 (1D3) (CL647-65290, Clone: 1D3), and 0.25 ug FITC Plus Anti-Mouse CD48 Rabbit RecAb (FITC-98081, Clone: 240732A5) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9). Cells were not fixed.



1x10^6 mouse splenocytes were surface stained with 0.25 ug FITC Plus Anti-Mouse CD48 Rabbit RecAb (FITC-98081, Clone: 240732A5) (red) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9) (blue). Cells were not fixed.