For Research Use Only

FITC Plus Anti-Human CD97 Rabbit Recombinant Antibody

www.ptglab.com

Purification Method:

Protein A purification

Recommended Dilutions:

FC: 5 ul per 10^6 cells in a 100 µl

Excitation/Emission maxima

CloneNo.:

241335D8

suspension

wavelengths:

495 nm / 524 nm

Catalog Number:FITC-98094-2

Basic Information

Catalog Number:

FITC-98094-2

Size:

100tests, 5 ul/test

Source: Rabbit

Isotype:

GenBank Accession Number:

BC026690 GeneID (NCBI):

UNIPROT ID: P48960

Full Name:

CD97 molecule Calculated MW:

722 aa, 79 kDa

Positive Controls: FC: human PBMCs,

Applications

Tested Applications:

Species Specificity:

human

Background Information

CD97 is a member of the seven-trans-membrane subfamily of the class B G protein-coupled receptor (GPCR) group of the epidermal growth factor (EGF) and is present on the surface of lymphocytes, monocytes, macrophages, dendritic cells, granulocytes, and smooth muscle, being a dimeric glycoprotein with a 75-90 kDa intracellular domain and a 28 kDa extracellular domain (PMID: 23969601). It plays a vital role in cell adhesion, migration, and cell connection regulation (PMID: 36464675). CD97 is expressed on T-cells, but rarely on B cells. CD97 might play key roles in cellular adhesion, through connections with other proteins of the cell surface and of the extracellular matrix (PMID: 32781778).

Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data





1x10^6 human PBMCs were surface stained with APC Anti-Human CD3 (OKT3) Mouse IgG2a Recombinant Antibody (APC-65569, Clone: OKT3), and 5 ul FITC Plus Anti-Human CD97 Rabbit RecAb (FITC-98094-2, Clone: 241335D8) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 human PBMCs were surface stained with 5 ul FITC Plus Anti-Human CD97 Rabbit RecAb (FITC-98094-2, Clone: 241335D8) (red) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9)(blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.