For Research Use Only

## FITC Plus Anti-Human CD7 Rabbit Recombinant Antibody

Catalog Number: FITC-98204



**Basic Information** 

Catalog Number: GenBank Accession Number: FITC-98204 NM 006137.7

GeneID (NCBI): Size:

100tests, 5 ul/test

ENSEMBL Gene ID: Source: Rabbit ENSG00000173762

UNIPROT ID: Isotype: P09564 Full Name: Immunogen Catalog Number:

EG3034 CD7 molecule

> Calculated MW: 25 kDa

**Purification Method:** 

Protein A purification

CloneNo.: 242402H8

Recommended Dilutions:

FC: 5 ul per 10^6 cells in a 100 µl

suspension

Excitation/Emission maxima

wavelengths: 495 nm / 524 nm

**Applications** 

**Tested Applications:** 

Species Specificity:

human

Positive Controls:

FC: human PBMCs,

## **Background Information**

CD7, also known as Leu-9 or GP40, is a 40-kDa single-pass type I transmembrane glycoprotein belonging to the immunoglobulin superfamily. CD7 is expressed on thymocytes, T cells, NK cells, and cells in the early stages of T, B, and myeloid cell differentiation (PMID: 10530432). It is one of the earliest antigens to appear on cells of the Tlymphocyte lineage (PMID: 3501369). CD7 plays a significant role in T-cell and T-cell/B-cell interactions during early lymphoid development. It is involved in the regulation of cell signaling and biology, including T-cell activation, proliferation, and the expression of interleukin-2 receptor alpha (IL-2Ra) (PMID: 11485208). CD7 is not only a marker for T-cells but also has functional implications in immune cell interactions and adhesion. It can provide co-stimulatory signals and is associated with CD3 and CD45, participating in T-cell activation (PMID: 7523512).

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 0.09% sodium azide, pH7.3

## **Selected Validation Data**





1x10^6 human PBMCs were surface stained with APC Anti-Human CD4 (OKT4) (APC-65134, Clone: OKT4), and 5ul FITC Plus Anti-Human CD7 Rabbit RecAb (FITC-98204, Clone: 242402H8) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 human PBMCs were surface stained with 5 ul FITC Plus Anti-Human CD7 Rabbit RecAb (FITC-98204, Clone: 242402H8) (red) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9) (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.