## For Research Use Only

## MultiPro<sup>TM</sup> 5CFLX Anti-Human Lselectin/CD62L (Polyclonal)




Catalog Number: G26477-1-5C


Basic Information	Catalog Number: G26477-1-5C	GenBank Accession Number: BC020758	Conjugate: 5CFLX				
	Size: 10ug, Concentration: 500ug/mL by Bradford method using BSA as the standard; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG24063	GeneID (NCBI): 6402 ENSEMBL Gene ID: ENSG00000188404 UNIPROT ID: P14151 Full Name: MultiPro <sup>TM</sup> 5CFLX Anti-Human L- selectin/CD62L (Polyclonal)	Full Oligo Sequence: CGGAGATGTGTATAAGAGACAGAGCT TCGCTGAACCGCCCATATAAGAAA Barcode Sequence: AGCTTCGCTGAACCG				
				Applications	Tested Applications: Single Cell (Intra)		
					Species Specificity: Human		
				Background Information	CD62L, also known as L-selectin or SELL, is a member of the selectin family of adhesion molecules that also include CD62E (E-selectin) and CD62P (P-selectin) (PMID: 2663882, 2473156, 1382078). CD62L is a highly glycosylated protein of 95-105 kDa on neutrophils and 74 kDa on lymphocytes (PMID: 1382078; 1694883, 1695155). CD62L is expressed on the surface of most leukocytes, including lymphocytes, neutrophils, monocytes, eosinophils, hematopoietic progenitor cells, and immature thymocytes (PMID: 1694883, 1688580). It mediates the binding of lymphocytes to high endothelial venules (HEV) of peripheral lymph nodes through interactions with a constitutively expressed ligand, and is also involved in lymphocyte, neutrophil, and monocyte attachment to endothelium at sites of inflammation (PMID: 1382078).		
Storage	Storage: 2-8°C Storage Buffer: PBS with 1mM EDTA and 0.09% sodi	um azide					

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





G26477-1-5C was used to stain Resting PBMC and analyzed with 10x Genomics Gene Expression Flex with Feature Barcodes and Multiplexing kit with Fix-Stain protocol. G26477-1-5C was used to stain PBMC under 4hr LPS + TI treatment and analyzed with 10x Genomics Gene Expression Flex with Feature Barcodes and Multiplexing kit with Fix-Stain protocol.