

HRP-conjugated GAPDH Monoclonal antibody

Catalog Number: HRP-60004

553 Publications

Basic Information

Catalog Number: HRP-60004	GenBank Accession Number: BC004109	Purification Method: Antigen affinity purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 2597	CloneNo.: 1E6D9
Source: Mouse	Full Name: glyceraldehyde-3-phosphate dehydrogenase	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG2b	Calculated MW: 36 kDa	
	Observed MW: 36 kDa	

Applications

Tested Applications: WB	Positive Controls: WB : HeLa cells, HepG2 cells, HEK-293 cells
Cited Applications: IF, IHC, IP, WB	
Species Specificity: human, mouse, rat, zebrafish, plant	
Cited Species: human, Chicken, rat, Honey bees, mouse, rabbit, hamster, pig, opossum	

Background Information

Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) catalyzes the phosphorylation of glyceraldehyde-3-phosphate during glycolysis. GAPDH participates in nuclear events including transcription, binding RNA, RNA transportation, DNA replication, DNA repair and apoptosis. Being stably and constitutively expressed at high levels in most tissues and cells, GAPDH is considered a housekeeping protein. It is widely used as a control for RT-PCR and also loading control in electrophoresis and Western blotting. GAPDH is normally expressed in cellular cytoplasm or membrane, but can occasionally translocate to the nucleus after the addition of post-translational modifications such as S-nitrosylation. This antibody is raised against full length GAPDH of human origin. It can recognize the 36 kDa GAPDH protein in most cells/tissues. In addition, a band below 36 kDa can always be detected as the isoform or spliced product of GAPDH (PMID: 23885286, 23877755, 19368702). Please note that some physiological factors, such as hypoxia and diabetes, increase GAPDH expression in certain cell types.

Notable Publications

Author	Pubmed ID	Journal	Application
Dong Yu	36183792	Brain Res	WB
Jiawei Hao	36126167	Autophagy	WB
Victoria L. Messerschmidt	34651022	Front Cardiovasc Med	WB

Storage


Storage:
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Western blot analysis of GAPDH in various tissues and cell lines using Proteintech antibody HRP-60004 at a dilution of 1:10000.