For Research Use Only

HRP-conjugated 6*His, His-Tag Monoclonal antibody Catalog Number:HRP-66005 166 Publications




Basic Information	Catalog Number: HRP-66005	GenBank Access	ion Number:	Purification Method: Protein G purification
	Size:	GeneID (NCBI): Full Name: Calculated MW:		CloneNo.: 187G5
	100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Mouse			
		0.84 kDa		Recommended Dilutions: WB 1:5000-1:50000
	Isotype: IgG1			
Applications	Tested Applications: WB	Positive Controls:		
	WB : Recombinant protein protein, Cited Applications: WB, IF, IP, CoIP, ELISA, Blocking assay			
	Species Specificity: recombinant protein			
Background Information	Protein tags are protein or peptide sequences located either on the C- or N- terminal of the target protein, which facilitates one or several of the following characteristics: solubility, detection, purification, localization and expression. His-tag is often used for affinity purification and binding assays. Expressed His-tagged proteins can be purified and detected easily because the string of histidine residues binds to several types of immobilized metal ions, including nickel, cobalt and copper, under specific buffer conditions. This His-tag antibody is a useful tool for monitoring of the His-tagged proteins, and recognizes His-tags placed at N-terminal, C-terminal, and internal regions of fusion proteins expressed in bacteria, insect, and mammalian cells. This antibody is conjugated with HRF Note: Do not add Azium (Sodium Azide or Smite) into the dilution buffer. Azium is the HRP inhibitor which decreases the enzymatic activity of HRP.			
	-	de or Smite) into t	ne dilution buffer. A	zium is the HRP inhibitor which decrease
Notable Publications	the enzymatic activity of HRP.			
Notable Publications	the enzymatic activity of HRP. Author P	de or Smite) into t ubmed ID 6178799	Journal	zium is the HRP inhibitor which decrease Application WB
Notable Publications	the enzymatic activity of HRP. Author P Antti Kulmala 3	ubmed ID		Application
Notable Publications	the enzymatic activity of HRP. Author P Antti Kulmala 3 Rudolf Hergesheimer 3	ubmed ID 6178799	Journal ACS Synth Biol	Application

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Recombinant protein were subjected to SDS PAGE followed by western blot with HRP-66005 (6°His, His-Tag antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.