For Research Use Only

HRP-conjugated Alpha Tubulin Monoclonal antibody



Catalog Number: HRP-66031

Featured Product

149 Publications

Basic Information

Catalog Number:

Size:

GenBank Accession Number:

HRP-66031 BC009314

GeneID (NCBI):

100ul , Concentration: 1000 ug/ml by 10376

UNIPROT ID:

Nanodrop: Mouse

P68363 Full Name:

Isotype: lgG2b

tubulin, alpha 1b Calculated MW:

50 kDa

Observed MW: 52 kDa

Applications

Tested Applications:

WB

WB, CoIP

Species Specificity:

human, mouse, rat

Purification Method:

Antigen affinity purification

CloneNo.: 1E4C11

Recommended Dilutions:

WB: 1:5000-1:50000

Cited Applications:

human, mouse, rat **Cited Species:**

Positive Controls:

WB: HeLa cells, HEK-293 cells

Background Information

There are five tubulins in human cells: alpha, beta, gamma, delta, and epsilon. Tubulins are conserved across species. They form heterodimers, which multimerize to form a microtubule filament. An alpha and beta tubulin heterodimer is the basic structural unit of microtubules. The heterodimer does not come apart once formed. The alpha and beta tubulins, which are each about 55 kDa MW, are homologous but not identical. Alpha tubulin is useful for scientists across fields as an internal control due to its high, ubiquitous expression pattern. Tubulin expression may vary according to resistance to antimicrobial and antimitotic drugs. This antiody recognize Tubulin-Alpha only and it is conjugated with HRP.

Note: Do not add Azium (Sodium Azide or Smite) into the dilution buffer. Azium is the HRP inhibitor which decreases the enzymatic activity of HRP.

Notable Publications

Author	Pubmed ID	Journal	Application
Immacolata Giordano	34621739	Front Cell Dev Biol	WB
Jiaxi Song	36125175	J Clin Endocrinol Metab	WB
Ruzhi Wei	30278376	Stem Cell Res	WB

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with HRP-66031 (Alpha Tubulin antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Western blot analysis of alpha-tubulin in various tissues and cell lines using Proteintech antibody HRP-60031 at a dilution of 1:5000.