For Research Use Only

## HRP-conjugated Beta Actin Recombinant antibody

Catalog Number: HRP-81115



**Basic Information** 

Catalog Number: GenBank Accession Number:

HRP-81115 BC002409
Size: GeneID (NCBI):

100ul , Concentration: 1000 ug/ml by 60 Nanodrop; UNIPROT ID:

Source: P60709
Rabbit Full Name:
Isotype: actin, beta
IgG Calculated MW:
Immunogen Catalog Number: 375 aa, 42 kDa
AG14521 Observed MW:
42 kDa

Protein A purification CloneNo.: 4H1

**Purification Method:** 

Recommended Dilutions: WB 1:2000-1:16000

**Applications** 

**Tested Applications:** 

W/R

Species Specificity: human, mouse, rat, pig

**Positive Controls:** 

WB: HeLa cells, rat brain tissue, mouse brain tissue

## **Background Information**

Beta Actin, also named as ACTB and F-Actin, belongs to the actin family. Actins are highly conserved globular proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells. At least six isoforms of actins are known in mammals and other vertebrates: alpha (ACTC1, cardiac muscle 1), alpha 1 (ACTA1, skeletal muscle) and 2 (ACTA2, aortic smooth muscle), beta (ACTB), gamma 1 (ACTG1) and 2 (ACTG2, enteric smooth muscle). Beta and gamma 1 are two non-muscle actin proteins. Most actins consist of 376aa, while ACTG2 (rich in muscles) has 375aa and ACTG1(found in non-muscle cells) has only 374aa. Beta actin has been widely used as the internal control in RT-PCR and Western Blotting as a 42-kDa protein. However, the 37-40, 31, 15 kDa cleaved fragment of beta actin can be generated during apoptosis process. This antibody was generated against N-terminal region of human beta actin protein and can cross-react with other actins. (9173887, 11217076, 10229193)

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with HRP-81115 (Beta Actin antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.