

Catalog Number: HZ-1138

Data Sheet



HumanKine® Activin A (Recombinant Human)

Animal Component-Free


Human cell expressed

Tag-Free


Endotoxin Free

Product Description				
Animal-free Recombinant Human Activin A is a member of the TGF beta superfamily, expressed from human 293 cells as disulfide linked homodimer with an apparent molecular mass of 25 kDa. This cytokine is produced in a serum-free, chemically defined media.				
Alternative Names	Activin A, Activin beta A chain, EDF, FRP, INHBA, Inhibin beta A chain, inhibin, beta A			
Source	Source Human Embryonic Kidney cells (HEK293). HEK293-derived Activin A protein			
Species Reactivity	human,mouse			

Specifications					
Test	Method	Specification			
Activity	Dose-dependent inhibition of the proliferation of the MPC-11 cell line (mouse plasmocytoma cell line)	0.5-3.5 ng/mL EC50			
Molecular Mass	SDS-PAGE	22 kDa, homodimer, non- glycosylated			
Purity	SDS-PAGE	>95%			
Endotoxin	LAL	<1 EU/µg			



Recombinant human Activin A (HZ-1138) causes dosedependent inhibition of proliferation in the Mouse plasmacytoma (MPC-11) cell line. Proliferation of the MPC-11 cell line was assessed using Promega CellTiter 96®. MPC-11 cells were treated with increasing concentrations of recombinant human Activin A for 72 hrs hours before addition of CellTiter96® reagent. The



www.ptglab.com

Document #: FR-QA118-101 Rev 0
Data Sheet Version #: 1

Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com

Preparation				
Shipping Temperature ambient temperature				
Formulation	1x PBS, See Certificate of Analysis for details			
Reconstitution	Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein in sterile 1xPBS pH 7.4 containing 0.1% endotoxin-free recombinant human serum albumin (HSA).			

Stability and Storage	Product Form	Temperature Conditions	Storage Time (From Date of Receipt)
	Lyophilized	-20°C to -80°C	Until Expiry Date
	Lyophilized	Room Temperature	2 weeks
	Reconstituted as per CofA	-20°C to -80°C	6 months
	Reconstituted as per CofA	4°C	1 week
		Avoid repeated freeze-thaw cycles.	

www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1 Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com