

Catalog Number: HZ-1204

Data Sheet



HumanKine® recombinant human VEGF121 protein

Animal Component-Free

Human cell expressed

Tag-Free


Endotoxin Free

Product Description


Animal-free Recombinant Human VEGF121 is expressed in human 293 cells as a glycosylated cytokine with an apparent molecular mass of 28 kDa as dimer and 40 kDa as trimer. Production in human 293 cells offers authentic glycosylation, contributing to stability in cell growth media. The cytokine is produced in a human cell expression system with serum free, chemically defined media. VEGF121 is a potent growth and angiogenic cytokine. It stimulates proliferation and survival of endothelial cells, and promotes angiogenesis and vascular permeability.

Alternative Names	nes L VEGFA, MVCD1, Vascular permeability factor, VEGF, VEGF A, VEGF121, VEGFA, VPF	
Source Human Embryonic Kidney cells (HEK293). HEK293-derived VEGF121 protein		
Species Reactivity	human,pig	

Specifications					
Test	Method	Specification			
Activity	Dose-dependent stimulation of the proliferation of HUVEC cells (Human Umbilical Vein Endothelial Cells)	1.5-9 ng/mL in Reporter cells, ≤ 15 ng/mL HUVEC cells			
Molecular Mass	SDS-PAGE	17 and 21 kDa reduced, 28 to 40 kDa non-reduced, homodimer and homotrimer, glycosylated			
Purity	SDS-PAGE	>95%			
Endotoxin	LAL	<1 EU/μg			



Recombinant human
VEGF121 (HZ-1204) induces
dose-dependent
proliferation of the HUVEC
(human umbilical vein
endothelial) cell line. Cell
number was quantitatively
assessed by PrestoBlue®
cell viability reagent. HUVEC
cells were treated with
increasing concentrations of
recombinant VEGF121 for 96
hours. The EC50 was
determined using a 4parameter non-linear



www.ptglab.com

Document #: FR-QA118-101 Rev 0
Data Sheet Version #: 1

Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com

Preparation				
Shipping Temperature	ambient temperature			
Formulation 1x PBS, See Certificate of Analysis for details				
Reconstitution	Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein to 0.2 mg/mL in sterile 1xPBS pH 7.4 containing 0.1% endotoxin-free recombinant human serum albumin (HSA). Gently swirl or tap vial to mix.			

Stability and Storage	Product Form	Temperature Conditions	Storage Time (From Date of Receipt)
	Lyophilized	-20°C to -80°C	Until Expiry Date
	Lyophilized	Room Temperature	2 weeks
	Reconstituted as per CofA	-20°C to -80°C	6 months
	Reconstituted as per CofA	4°C	1 week
		Avoid repeated freeze-thaw cycles.	

www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1 Proteintech Group, Inc. 5500 Pearl Street, Suite 400 Rosemont, IL 60612

t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com