

Catalog Number: HZ-1308

Data Sheet



HumanKine® recombinant human PDGFbb protein

Animal Component-Free

Human cell expressed

Tag-Free


Endotoxin Free

Product Description


Animal-free Recombinant Human PDGFbb is a growth factor that promotes blood vessel formation, mitogenesis, chemotaxis, etc. The PDGF family members (four homodimers and one heterodimer) are secreted, disulfide-linked dimeric glycoproteins that regulate their cellular functions through interaction with PDGFR receptors. PDGFbb is synthesized, stored, and released by alpha granules of platelets. It is also known as PDGF-2, becaplermin, or GDGF. Dysregulation in PDGF signaling has been shown to be associated with tumorigenesis and progression of cancer. Recombinant PDGFbb is used in treatment of chronic ulcers and to speed healing in surgical procedures.

Alternative Names	Becaplermin, c sis, FLJ12858, PDGF 2, PDGF subunit B, PDGF2, PDGFB, PDGFbb, Proto oncogene c Sis, SIS, SSV		
Source	Human Embryonic Kidney cells (HEK293). HEK293-derived PDGFbb protein		
Species Reactivity	human,mouse,rat		

Specifications					
Test	Method	Specification			
Activity	Dose-dependent stimulation of the proliferation of 3T3 cells	0.3-3 ng/mL			
Molecular Mass	SDS-PAGE	14 and 36 kDa reduced, 29 to 32 kDa non-reduced, homodimer, glycosylated			
Purity	SDS-PAGE	>95%			
Endotoxin	LAL	<1 EU/µg			



Recombinant human PDGFbb (HZ-1308) stimulates dose-dependent proliferation of the NIH/3T3 mouse fibroblast cell line. Viable cell number was quantitatively assessed by Prestoblue Cell Viability Reagent. NIH/3T3 cells were serum starved with 0.1% FBS for 24 hours before treatment with increasing concentrations of recombinant human PDGFbb for 48 hrs. The EC50 was



www.ptglab.com

Document #: FR-QA118-101 Rev 0
Data Sheet Version #: 1

Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com

Preparation				
Shipping Temperature ambient temperature				
Formulation	Formulation 50mM Acetate pH 3.5 + 500mM NaCl, See Certificate of Analysis for details			
Reconstitution Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein to 0.2 mg/mL in sterile 1x containing 0.1% endotoxin-free recombinant human serum albumin (HSA). Gently swirl or tap vial to mix.				

	Product Form	Temperature Conditions	Storage Time (From Date of Receipt)
	Lyophilized	-20°C to -80°C	Until Expiry Date
Stability and Storage	Lyophilized	Room Temperature	2 weeks
	Reconstituted as per CofA	-20°C to -80°C	6 months
	Reconstituted as per CofA	4°C	1 week
		Avoid repeated freeze-thaw cycles.	

www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1 Proteintech Group, Inc. 5500 Pearl Street, Suite 400 Rosemont, IL 60612

t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com