



IHCeasy® CD22 Ready-To-Use IHC Kit

Catalog Number: KHC0019

General Information

Sample type: FFPE tissue Cited sample type: Reactivity: Human Cited Reactivity: Assay type: Immunohistochemistry Primary antibody type: Mouse Monoclonal

Secondary antibody type: Polymer-HRP-Goat anti-Mouse

Kit Component

Component	Size	Concentration
Antigen Retrieval Buffer	100 mL	50×
Washing Buffer	100 mL ×2	20×
Blocking Buffer	5 mL	RTU
Primary Antibody	5 mL	RTU
Secondary Antibody	5 mL	RTU
Chromogen Component A	0.2 mL	RTU
Chromogen Component B	4 mL	RTU
Signal Enhancer	5 mL	RTU
Counter Staining Reagent	5 mL	RTU
Mounting Media	5 mL	RTU
Control Slide	1 slide (Optional)	FFPE
Datasheet	1 Copy	
Manual	1 Copy	

Storage Instructions

All the reagents are stored at 2-8°C. The kit is stable for 6 months from the date of receipt.


Background

CD22, also known as Siglec-2 (sialic acid binding Ig-like lectin 2) or BL-CAM (B-lymphocyte cell adhesion molecule), is a 130-140 kDa, B-cell restricted, type I transmembrane glycoprotein belonging to the immunoglobulin gene superfamily. The expression of CD22 is developmentally regulated. It is expressed at low levels in the cytoplasm of pro-B and pre-B cells and present on the cell surface only at mature stages of B-cell differentiation. Cell surface expression is lost during terminal differentiation into plasma cell and after B-cell activation. CD22 is an inhibitory receptor for B-cell receptor (BCR) signalling, preferentially binds to alpha-2,6-linked sialic acid and mediates B-cell B-cell interactions. It plays a crucial role in activation and differentiation of the B-cell.


Synonyms

B cell receptor CD22, BL CAM, CD22, CD22 molecule, SIGLEC 2, SIGLEC2, T cell surface antigen Leu 14

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using KHC0019 (CD22 IHC Kit).



Immunohistochemical analysis of paraffinembedded human appendicitis tissue slide using KHC0019 (CD22 IHC Kit).