



IHCeasy TTF2 Ready-To-Use IHC Kit

Catalog Number: KHC2007

General Information

Sample type: FFPE tissue Cited sample type: Reactivity: Human, Mouse, Rat Cited Reactivity: Assay type: Immunohistochemistry Primary antibody type: Rabbit Polyclonal

Secondary antibody type: Polymer-HRP-Goat anti-Rabbit

Kit Component

Antigen Retrieval Buffer 100 mL	50×
Washing Buffer 100 mL × 2	20×
Blocking Buffer 5 mL	RTU
Primary Antibody 5 mL	RTU
Secondary Antibody 5 mL	RTU
Chromogen Component A 0.2 mL	RTU
Chromogen Component B 4 mL	RTU
Signal Enhancer 5 mL	RTU
Counter Staining Reagent 5 mL	RTU
Mounting Media 5 mL	RTU
Control Slide 1 slide (Optional)	FFPE
Datasheet 1 Copy	
Manual 1 Copy	

Storage Instructions

All the reagents are stored at 2-8°C. The kit is stable for 6 months from the date of receipt.


Background

Transcription elongation is an key target for control of eukaryotic gene expression. After initiating from a promoter, the RNA polymerase II elongation complex is subjected to a block to elongation by the action of components of N-TEF (negative transcription elongation factor). Transcription termination factor 2 (TTF2) is a DsDNA-dependent ATPase that acts as a transcription termination factor by coupling ATP hydrolysis with removal of RNA polymerase II from the DNA template. It also contribute to mitotic transcription repression, and involved in pre-mRNA splicing.


Synonyms

F2, HuF2, Lodestar homolog, Transcription release factor 2, TTF2


Selected Validation Data



Immunohistochemical analysis of paraffinembedded mouse lung tissue slide using KHC2007 (TTF2 IHC Kit).



Immunohistochemical analysis of paraffinembedded mouse stomach tissue slide using KHC2007 (TTF2 IHC Kit).



Immunohistochemical analysis of paraffinembedded rat stomach tissue slide using KHC2007 (TTF2 IHC Kit).