For Research Use Only

NGAL Recombinant Matched Antibody Pair, PBS Only



Catalog Number: MP00770-1

Capture Antibody Information

Catalog Number: 82841-9-PBS Host:

Rabbit Isotype:

Purification Method: Protein A purification Clone ID: 2N10

Reactivity: human

Conjugate: Unconjugated Full name: lipocalin 2 Gene ID: 3934

Detection Antibody Information

Catalog Number: 82841-2-PBS Rabbit Isotype:

IgG **Purification Method:** Protein A purification

Clone ID: 218 Reactivity:

human GenBank: NP_005555.2 Conjugate: Unconjugated

Full name: lipocalin 2 Gene ID: 3934

Applications

Tested Applications: Sandwich ELISA

19.5-1250 pg/mL (Sandwich ELISA)

Recommended Dilutions:

It is recommended that this reagent should be titrated in each testing system to obtain optimal results.

Product Information

MP00770-1 targets NGAL in immunoassays as a matched antibody pair. Validated in Sandwich ELISA.

Capture antibody: Lipocalin-2/NGAL Recombinant antibody, PBS Only (Capture) 82841-9-PBS (2N10). 100 µg. Concentration 1 mg/ml.

 $Detection\ antibody:\ Lipocalin-2/NGAL\ Recombinant\ antibody,\ PBS\ Only\ (Detector)\ 82841-2-PBS\ (2L8).\ 100\ \mu g.$ Concentration 1 mg/ml.

Unconjugated rabbit recombinant monoclonal antibody pair in PBS only storage buffer at a concentration of 1 mg/mL, ready for conjugation. Created using Proteintech's proprietary in-house recombinant technology. Recombinant production enables unrivalled batch-to-batch consistency, easy scale-up, and future security of supply.

Matched antibody pairs are designed for use in a variety of assays and platforms that require matched antibody


Antibody use should be optimized for each application and assay.


Storage


Storage:

Store at -80°C. Storage buffer: PBS only


Selected Validation Data









Plasma of fifteen individual healthy human donors was measured. The NGAL concentration of detected samples was determined to be 38,260.8 pg/mL with a range of 17,778.7 - 90,593.8 pg/mL



Human peripheral blood mononuclear cells (PBMC) were cultured unstimulated or stimulated with 10 µg/mL PHA for 1 days. The mean NGAL concentration was determined to be 18,080.8 pg/mL in unstimulated PBMC supernatant, 32,268.8 pg/mL in PHA stimulated PBMC supernatant.



Saliva of eight individual healthy human donors was measured. The NGAL concentration of detected samples was determined to be 158,732.5 pg/mL with a range of 52,136.9 - 388,400.2 pg/mL



Urine of eight individual healthy human donors was measured. The NGAL concentration of detected samples was determined to be 11,654.6 pg/mL with a range of 1,964.2 - 21,150.4 pg/mL