For Research Use Only

IL-16 Recombinant Matched Antibody Pair, PBS Only

www.ptglab.com

Catalog Number: MP02772-1

Capture Antibody Information

Catalog Number: Clone ID: 86927-2-PBS 251661G11 Host: Reactivity: Rabbit human

Isotype Immunogen Catalog Number: Eg4831

Purification Method: Protein A purification Unconjugated Full name:

Conjugate:

interleukin 16 (lymphocyte chemoattractant factor)

Gene ID: 3603

Detection Antibody Information

Catalog Number: Clone ID: Conjugate: 86927-1-PBS 251661B4 Host: Reactivity: Full name: Rabbit human

Isotype: GenBank: IgG AAC12732.1

Purification Method: Immunogen Catalog Number:

Protein A purification Eg4831 Unconjugated

interleukin 16 (lymphocyte chemoattractant factor)

Gene ID: 3603

Applications

Tested Applications:

0.781-100 ng/mL (Cytometric Bead Cytometric bead array

Array)

Recommended Dilutions:

It is recommended that this reagent should be titrated in each testing system to obtain optimal results.

Product Information

MP02772-1 targets IL-16 in immunoassays as a matched antibody pair. Validated in Cytometric bead array.

Capture antibody: IL-16 Recombinant monoclonal antibody, PBS Only (Capture) 86927-2-PBS (251661G11). 100 µg.

 $Detection\ antibody:\ LI-16\ Recombinant\ monoclonal\ antibody,\ PBS\ Only\ (Detector)\ 86927-1-PBS\ (251661B4).\ 100\ \mu g.$ Concentration 1 mg/ml.

Unconjugated rabbit recombinant monoclonal antibody pair in PBS only storage buffer at a concentration of 1 mg/mL, ready for conjugation. Created using Proteintech's proprietary in-house recombinant technology. Recombinant production enables unrivalled batch-to-batch consistency, easy scale-up, and future security of supply.

Matched antibody pairs are designed for use in a variety of assays and platforms that require matched antibody


Antibody use should be optimized for each application and assay.


Storage


Storage:

Store at -80°C. Storage buffer: PBS only

Selected Validation Data







Human peripheral blood mononuclear cells (PBMC) were cultured unstimulated or stimulated with 10 µg/mL PHA for 5 days. The mean IL-16 concentration was determined to be 5.5 ng/mL in unstimulated PBMC supernatant, 15.5 ng/mL in PHA stimulated PBMC supernatant.