For Research Use Only

## IL17A Monoclonal Matched Antibody Pair, PBS Only



Catalog Number: MP50004-1

**Capture Antibody** Information

Catalog Number: 69021-2-PBS Host:

Mouse

Isotype: lgG1

**Purification Method:** Protein G purification

**Detection Antibody** Information

Catalog Number: 69021-1-PBS Mouse Isotype: lgG1

**Purification Method:** 

Clone ID: 1E3H1 Reactivity:

human Immunogen Catalog Number: HZ-1113

Conjugate:

Conjugate:

Full name:

Gene ID:

3605

Unconjugated

interleukin 17A

Unconjugated Full name: interleukin 17A Gene ID:

3605

Protein G purification

**Applications** 

**Tested Applications:** Sandwich ELISA

Clone ID:

Reactivity:

GenBank:

HZ-1113

1F3E3

human

Immunogen Catalog Number:

Recommended Dilutions:

31.25-2000 pg/mL (Sandwich ELISA) It is recommended that this reagent should be titrated in each testing system to obtain optimal results.

**Product Information** 

MP50004-1 targets IL17A in immunoassays as a matched antibody pair. Validated in Sandwich ELISA.

Capture antibody: IL-17A Monoclonal antibody, PBS Only (Capture) 69021-2-PBS (1E3H1). 100 µg. Concentration 1

 $Detection\ antibody;\ Neutra Kine @\ IL-17A\ Monoclonal\ antibody,\ PBS\ Only\ (Detector)\ 69021-1-PBS\ (1F3E3).\ 100\ \mu g.$ Concentration 1 mg/ml.

Unconjugated mouse monoclonal antibody pair in PBS only storage buffer at a concentration of 1 mg/mL, ready for conjugation.

Matched antibody pairs are designed for use in a variety of assays and platforms that require matched antibody


Antibody use should be optimized for each application and assay.

Storage


Storage: Store at -80°C. Storage buffer:

PBS only


## **Selected Validation Data**



Sandwich ELISA standard curve of MP50004-1, IL17A Monoclonal Matched Antibody Pair, PBS Only. Capture antibody: 69021-2-PBS. Detection antibody: 69021-1-PBS. Standard: HZ-1113. Range: 31.25-2000 pg/mL



Twenty-four serum samples from volunteers were evaluated for human IL-17A in this assay. twenty-two samples measured less than the lowest standard, 31.25 pg/mL Two samples measured 98.5 pg/mL and 107.5 pg/mL respectively. No medical histories were available for the donors used in this study.



Human peripheral blood mononuclear cells (PBMC) were cultured unstimulated or stimulated with 10 µg/mL PHA for 3 days or 5 days. The mean IL17A concentration was undetected in unstimulated PBMC supernatant, 126.6 pg/mL in PBMC supernatant stimulated with PHA for 1 day and 251.8 pg/mL in PBMC supernatant stimulated with PHA for 5 days.