

For Research Use Only

FcZero-rAb™ PE-CoraLite® Plus 594 Anti-Human CD8 (SK1) Rabbit IgG Recombinant Antibody

Catalog Number:**PCL594-FcA65630**



Basic Information

Catalog Number:	PCL594-FcA65630	GenBank Accession Number:	BC025715	Purification Method:	Protein A purification
Size:	100 tests, 5 µl/test	GeneID (NCBI):	925	Clone No.:	SK1
Source:	Rabbit	ENSEMBL Gene ID:	ENSG00000153563	Recommended Dilutions:	FC: 5 µl per 10 ⁶ cells in a 100 µl suspension
Isotype:	IgG	UNIPROT ID:	P01732	Excitation/Emission maxima wavelengths:	496 nm, 565 nm / 604 nm
		Full Name:	CD8a molecule		
		Calculated MW:	235 aa, 26 kDa		

Applications

Tested Applications:	Positive Controls:
FC	FC: human PBMCs,
Species Specificity:	human, monkey, non-human primates

Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8 α or as a heterodimer of CD8 α and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 $\alpha\beta$ heterodimer while CD8 $\alpha\alpha$ homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) $\gamma\delta$ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).


Storage

Storage:
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 0.09% sodium azide and 0.5% BSA, pH7.3


For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ human PBMCs were surface stained with FcZero-rAb™ Coralite® Plus 750 Anti-Human CD3 (UCHT1), and 5 ul PE-CL594 Plus Anti-Human CD8 (SK1) Rabbit IgG RecAb (PCL594-FcA65630, Clone:SK1) or FcZero-rAb™ PE-Coralite® Plus 594 Rabbit IgG Isotype Control Recombinant Antibody (PCL594-FcA98136, Clone: 240953C9). Cells were not fixed. Lymphocytes were gated.



1x10⁶ human PBMCs were surface stained with 5 ul PE-CL594 Plus Anti-Human CD8 (SK1) Rabbit IgG RecAb (PCL594-FcA65630, Clone:SK1) (red) or FcZero-rAb™ PE-Coralite® Plus 594 Rabbit IgG Isotype Control Recombinant Antibody (PCL594-FcA98136, Clone: 240953C9) (blue). Cells were not fixed. Lymphocytes were gated.