## For Research Use Only

## PE Anti-Human CD11c (3.9)

Catalog Number: PE-65086



**Basic Information** 

Catalog Number:

GenBank Accession Number: BC038237

**Purification Method:** Affinity purification

PE-65086

GeneID (NCBI):

100tests , 5 µl/test

3687

CloneNo.: 3.9

Full Name:

Excitation/Emission maxima

Mouse

integrin, alpha X (complement component 3 receptor 4 subunit) wavelengths:

Isotype: IgG1, kappa

Calculated MW: 1169 aa, 129 kDa 496 nm, 565 nm / 578 nm

**Applications** 

**Tested Applications:** 

Species Specificity:

**Background Information** 


Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated  $\alpha$  and  $\beta$  subunits (PMID: 9779984). CD11c, also known as integrin aX, is a type I transmembrane glycoprotein present on a variety of cells, including monocytes/macrophages, granulocytes, a subset of B cells, NK cells and dendritic cells (PMID: 2897326; 1680915; 1694698; 17389580). As a result of its high level of expression on most dendritic cells, CD11c is typically considered to be a marker of conventional dendritic cells (PMID: 27119555). CD11c forms an  $\alpha/\beta$ heterodimer with CD18 (integrin  $\beta$ 2). CD11c/CD18 acts a receptor for fibrinogen and is important in monocyte adhesion and chemotaxis (PMID: 1671533).

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA.

## Selected Validation Data



1X10^6 human PBMCs were surface stained with 5 ul PE Anti-Human CD11c (PE-65086, Clone:3.9) or unstained. Cells were not fixed. Monocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul PE Anti-Human CD11c (PE-65086, Clone:3.9) or unstained. Cells were not fixed. Monocytes were gated