## For Research Use Only

## PE Anti-Human CD38 (HB7)

Catalog Number: PE-65107



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** Affinity purification

PE-65107 Size:

BC007964 GeneID (NCBI):

CloneNo.:

100tests , 5 µl/test

952

Excitation/Emission maxima

496 nm, 565 nm / 578 nm

Mouse

Full Name: CD38 molecule Calculated MW:

wavelengths:

Isotype: IgG1, kappa

300 aa, 34 kDa

**Applications** 

**Tested Applications:** 

Species Specificity:

**Background Information** 

 ${\tt CD38, also \, known \, as \, ADP-ribosyl \, cyclase \, 1, is \, a \, type \, II \, transmembrane \, glycoprotein \, with \, a \, short \, N-terminal \, and \, control \,$ cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four Nglycosylation sites (PMID: 2319135). The extracellular domain of CD38 has bifunctional enzyme activities that catalyze synthesis of cyclic ADP ribose from nicotinamide adenine dinucleotide (NAD) and hydrolysis of cyclic ADP ribose to adenosine diphosphoribose (PMID: 10636863). CD38 is expressed on a variety of hematopoietic and nonhematopoietic cells and is involved in diverse processes such as generation of calcium-mobilizing metabolites, cell activation, and chemotaxis (PMID: 25938500).


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Phosphate based buffer with 0.09% sodium azide and 0.1% gelatin, pH 7.2.

in USA), or 1(312) 455-8498 (outside USA)

## Selected Validation Data



1X10^6 human peripheral blood lymphocytes were surface stained with 5.00 ul/test PE Anti-Human CD38 (PE-65107, Clone:HB7) (red) or isotype control antibody (blue). Cells were not fixed.