For Research Use Only

PE Anti-Human CD163 (GHI/61)

Catalog Number: PE-65169 6 Publications



Basic Information

Catalog Number:

PE-65169

Size:

100tests , 5 µl/test

Source: Mouse

Isotype: IgG1, kappa GenBank Accession Number:

BC051281 GeneID (NCBI):

9332

ENSEMBL Gene ID:

ENSG00000177575 UNIPROT ID:

Full Name:

Q86VB7

CD163 molecule Calculated MW:

1156 aa, 125 kDa

Purification Method:

The purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. The conjugate is

purified by size-exclusion chromatography.

CloneNo.:

GHI/61

Recommended Dilutions:

FC: 5 ul per 10^6 cells in 100 µl

suspension

Excitation/Emission maxima

wavelengths:

496 nm, 565 nm / 578 nm

Applications

Tested Applications:

Cited Applications:

Species Specificity:

Human

Cited Species:

human

Positive Controls:

FC: human PBMCs.

Background Information

CD163, also known as M130, is a membrane glycoprotein which belongs to the scavenger receptor superfamily (PMID: 8370408). It is an acute phase-regulated and signal-inducing macrophage protein expressed exclusively in monocytes and tissue macrophages (PMID: 11196644). CD163 mediates endocytosis of haptoglobin-haemoglobin complexes (PMID: 11196644). The uptake of haptoglobin by macrophages contributes to the recycling of iron and also to the inflammatory response (PMID: 22900885). Soluble CD163 (sCD163), as a result of ectodomain shedding during inflammatory activation of macrophages, circulates in blood and has been suggested as a plasma/serum marker for macrophage activity (PMID: 12570164).

Notable Publications

Author	Pubmed ID	Journal	Application
Tao Yan	35410993	Cell Death Discov	FC
Yijun Lu	40184814	Int Immunopharmacol	FC
Tianci Han	39891717	Cancer Immunol Immunother	FC


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data



1X10^6 human PBMCs were treated with FC receptor block then surface stained with CL405 Anti-Human CD14 and 5 ul PE Anti-Human CD163 (PE-65169, Clone:GHI/61) or PE Mouse IgG1 Isotype Control. Cells were not fixed. Monocytes were gated.



1X10^6 human PBMCs were treated with FC receptor block then surface stained with 5 ul PE Anti-Human CD163 (PE-65169, Clone:GHI/61) or PE Mouse IgG1 Isotype Control. Cells were not fixed. Monocytes were gated.