For Research Use Only

PE Anti-Human CD141/Thrombomodulin Rabbit Recombinant Antibody

Size:



Catalog Number: PE-98037

Basic Information

Catalog Number: PE-98037

100tests, 5 ul/test Source:

Rabbit Isotype: IgG

Immunogen Catalog Number:

GenBank Accession Number:

NM 000361.3 GeneID (NCBI):

UNIPROT ID: P07204

7056

Full Name: thrombomodulin

Calculated MW: 60 kDa

Purification Method:

Protein A purification

CloneNo.: 240365H3

Recommended Dilutions:

FC: 5 ul per 10^6 cells in a 100 µl

suspension

Excitation/Emission maxima

wavelengths:

496 nm, 565 nm / 578 nm

Applications

Tested Applications:

Species Specificity:

human

Positive Controls:

FC: human peripheral blood leukocytes,

Background Information

Thrombomodulin, also known as CD141, is an endothelial cell surface glycoprotein that forms a 1:1 complex with the coagulation factor thrombin and plays an important role as a natural anticoagulant. Thrombomodulin serves to convert thrombin from a procoagulant protein into the activator for protein C. Once converted to activated protein C (APC), this protein serves as a major anticoagulant in blood (PMID: 2827310). Thrombomodulin is also located in $other cells \ (keratinocytes, osteoblasts, macrophages, \ldots) \ where \ it \ might be \ involved \ in \ cell \ differentiation \ or \ in \ differentiation \ or \ differentiation \ din \ differentiation \ differentiation \ differentiation \ differe$ inflammation (PMID: 9814688). In humans, thrombomodulin is encoded by the THBD gene. Mutations in this gene are a cause of thromboembolic disease, also known as inherited thrombophilia. Thrombomodulin is glycosylated and has an apparent molecular weight of 75 to 110 kDa (PMID: 1650405; 2827310).


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data



1x10^6 human peripheral blood leukocytes were surface stained with 5 ul PE Anti-Human CD141/Thrombomodulin Rabbit RecAb (PE-98037, Clone: 240365H3) or PE Rabbit IgG Isotype Control RecAb (PE-98136, Clone: 240953C9). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 FC receptor blocked human peripheral blood monocytes were surface stained with 5 ul PE Anti-Human CD141/Thrombomodulin Rabbit RecAb (PE-98037, Clone: 240365H3) (red) or PE Rabbit IgG Isotype Control RecAb (PE-98136, Clone: 240953C9) (blue). Cells were not fixed.