For Research Use Only

## FcZero-rAb™ PE Anti-Human CD79a (HM57) Rabbit IgG Recombinant Antibody



Catalog Number: PE-FcA65610

**Basic Information** 

Catalog Number: PE-FcA65610

Size:

100tests, 5 ul/test

Source: Rabbit

Isotype:

GenBank Accession Number:

BC113731 GeneID (NCBI):

Full Name:

CD79a molecule, immunoglobulin-

associated alpha

**Purification Method:** 

Protein A purification CloneNo.:

HM57

Excitation/Emission maxima

wavelengths:

496 nm, 565 nm / 578 nm

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity:

human

**Background Information** 

CD79a, also named as B-cell antigen receptor complex-associated protein alpha chain or MB-1 membrane glycoprotein, is a 226 amino acid protein, which contains one ITAM domain and one Ig-like C2-type (immunoglobulin-like) domain. CD79a is expressed in B cell and localizes in the cell membrane. CD79a is required in cooperation with CD79b for initiation of the signal transduction cascade activated by binding of antigen to the Bcell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. CD79a is also required for BCR surface expression and for efficient differentiation of proand pre-B-cells.

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer

PBS with 0.09% sodium azide and 0.5% BSA.

in USA), or 1(312) 455-8498 (outside USA)

## Selected Validation Data





1x10^6 human PBMCs were intracellularly stained with FcZero-rAb™APC Anti-Human CD19 (4G7), and 5 ul PE Anti-Human CD79a (HM57) Rabbit IgG RecAb (PE-FcA65610, Clone:HM57) or PE Rabbit IgG Isotype Control RecAb (PE-FcA98136, Clone: 240953C9). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).