## For Research Use Only

## FcZero-rAb<sup>™</sup> PE Anti-Human CD79a Rabbit Recombinant Antibody

Catalog Number: PE-FcA98072




| Basic Information      | Catalog Number:<br>PE-FcA98072  | GenBank Accession Number:<br>BC 113733  | Purification Method:<br>Protein A purification                        |
|------------------------|---|---|---|
|                        | Size:<br>100tests , 5 ul/test   | GenelD (NCBI):<br>973   | CloneNo.:<br>241299F3   |
|                        | Source:<br>Rabbit<br>Isotype:<br>IgG  | Full Name:<br>CD79a molecule, immunoglobulin-<br>associated alpha<br>Calculated MW:<br>226 aa, 25 kDa | Excitation/Emission maxima<br>wavelengths:<br>496 nm, 565 nm / 578 nm |
|                        |   |   |   |
| Background Information | CD79a, also named as B-cell antigen receptor complex-associated protein alpha chain or MB-1 membrane glycoprotein, is a 226 amino acid protein, which contains one ITAM domain and one Ig-like C2-type (immunoglobulin-like) domain. CD79A is expressed in B cells and localizes in the cell membrane. CD79A is required in cooperation with CD79B for initiation of the signal transduction cascade activated by binding of antigen to the B-cell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. CD79A is also required for BCR surface expression and for efficient differentiation of pro- and pre-B-cells. |   |   |
| Storage                | Storage:<br>Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.<br>Storage Buffer:<br>PBS with 0.09% sodium azide and 0.5% BSA, pH7.3  |   |   |

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1x10<sup>^</sup>6 human PBMCs were stained with APC Anti-Human CD19 (4G7), and 5 ul FcZero-rAb<sup>™</sup> PE Anti-Human CD79a Rabbit RecAb (PE-FcA98072, Clone:241299F3) or PE Rabbit IgG Isotype Control RecAb (PE-FcA98136, Clone: 240953C9). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).