For Research Use Only

FcZero-rAb™ PE-Cyanine5.5 Anti-Human CD18 (TS1/18) Rabbit IgG Recombinant Antibody



Purification Method:

CloneNo.:

suspension

wavelengths:

TS1/18

Protein A purification

Recommended Dilutions:

FC: 5 ul per 10^6 cells in a 100 µl

Excitation/Emission maxima

488 nm, 561 nm / 692 nm

Catalog Number: PY5P5-FcA65595

Basic Information

Catalog Number:

PY5P5-FcA65595

Size:

100tests, 5 ul/test

Source: Rabbit

Isotype: IgG

GenBank Accession Number:

GeneID (NCBI):

ENSEMBL Gene ID: ENSG00000160255

Full Name:

Calculated MW:

85 kDa

Applications

Tested Applications:

Species Specificity:

human

BC005861

3689

UNIPROT ID:

P05107

integrin, beta 2 (complement

component 3 receptor 3 and 4 subunit)

Positive Controls:

FC: human PBMCs,

Background Information

CD18, also known as integrin beta-2, is a transmembrane protein that forms heterodimers with CD11a, CD11b, CD11c, CD11d (PMID: 1968349). CD11a/CD18 (LFA-1) is expressed by all leukocytes, while CD11b/CD18 (Mac-1) and CD11c/CD18 (p150,95) are normally restricted to expression on monocytes, macrophages, polymorphonuclear lymphocytes (PMNs), and natural killer cells (PMID: 1968349; 3109455; 10946284). CD18 plays an important role in $mediating\ cell\ adhesion\ during\ immune\ response\ and\ defects\ of\ CD18\ cause\ leukocyte\ adhesion\ deficiency\ (PMID:$ 3594570).


Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data



1x10^6 human PBMCs were surface stained with CoraLite® Plus 405 Anti-Human CD45, and 5 ul PE-Cyanine5.5 Anti-Human CD18 (TS1/18) Rabbit IgG RecAb (PY5P5-FcA65595, Clone: TS1/18) or Rabbit IgG Isotype Control RecAb (PY5P5-FcA98136, Clone: 240953C9). Cells were not fixed.