

For Research Use Only

# FcZero-rAb® PE-Cyanine7 Anti-Rat CD8a (OX-8) Rabbit Recombinant Antibody

Catalog Number: PY7-FcA65577



## Basic Information

Catalog Number:

PY7-FcA65577

Size:

100ug , 200 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC088126

GeneID (NCBI):

24930

UNIPROT ID:

P07725

Full Name:

CD8a molecule

Purification Method:

Protein A purification

CloneNo.:

OX-8

Recommended Dilutions:

FC: 0.1 ug per 10<sup>6</sup> cells in a 100 µl suspension

Excitation/Emission maxima wavelengths:

496 nm, 565 nm / 778 nm

## Applications

Tested Applications:

FC

Species Specificity:

rat

Positive Controls:

FC : rat splenocytes,

## Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8aβ heterodimer while CD8aα homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) γδ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).

## Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1x10<sup>6</sup> rat splenocytes were surface stained with CoraLite® Plus 405 Anti-Rat CD4, and 0.1 ug PE-Cyanine7 Anti-Rat CD8a (OX-8) Rabbit RecAb (PY7-FcA65577, Clone: OX-8) or 0.1 ug Rabbit IgG Isotype Control RecAb (PY7-FcA98136, Clone: 240953C9). Cells were not fixed.



1x10<sup>6</sup> rat splenocytes were surface stained with 0.1 ug PE-Cyanine7 Anti-Rat CD8a (OX-8) Rabbit RecAb (PY7-FcA65577, Clone: OX-8) (red) or 0.1 ug Rabbit IgG Isotype Control RecAb (PY7-FcA98136, Clone: 240953C9) (blue). Cells were not fixed.